



ISSN: 0067-2904

## A Class of Fuzzy Integral Equations and the Existence and Uniqueness of Their Solutions

Alan Jalal Abdulqader and Sameer Qasim Hasan

Department of Mathematic, College of Education, Mustansiriyah university

Received: 4/10/2023

Accepted: 15/1/2024

Published: 30/1/2025

### Abstract

The aim of this paper to presented a class of interesting fuzzy integral nonlinear equations involving nonlinear fuzzy function of other multi fuzzy nonlinear functions that presented in fist time, the formula under study is an interesting type of a fuzzy nonlinear Volterra integral equation which defined on fuzzy numbers that has lower and upper functions. The Housdorf fuzzy metric space on fuzzy numbers and a fuzzy metric space on a fuzzy continuous functions and the space of all uniform modulus of absolutely fuzzy continuous functions have been given in details. Also the observer results investigated on parameter cases of fuzzy problem formulation, some necessary and sufficient conditions of boundedness and locally Lipschitz for different fuzzy functions and differential of them and also using Leibnitz rule for presented fuzzy problem formulation. The locally existence and uniqueness solutions in the space of all uniform modulus of absolutely fuzzy continuous functions which is generalized of fuzzy number space and established on contraction condition which proved by using the properties of fuzzy metric space this certain space. Also used concept of fuzzy Riemann integral and normed function to prove interesting results and illustrated in some examples to clear the proposal fuzzy integral nonlinear equations has a solution in certain fuzzy space that in firstly explained with different metrics spaces and normed properties and satisfied all the hypothesis for the component of the equation and obtain the estimations of the suggested conditions on certain fuzzy metrics as a stuffiest condition for grantee the assuming equations has a fuzzy solution with their lower and upper parameters.

**Keywords:** fuzzy number, nonlinear voltera integral, uniform modulus of absolutely continuous functions, fuzzy Riemann integral, existence and uniqueness

### صنف من المعادلات التكاملية الضبابية والوجودية والوحدانية لحلولها

الان جلال عبد القادر ، سمير قاسم حسن

قسم الرياضيات، كلية التربية، الجامعة المستنصرية، بغداد، العراق

### الخلاصة

الهدف من هذا البحث هو عرض صنف من المعادلات التكاملية الضبابية غير الخطية تحوي على دالة ضبابية غير خطية لدوال أخرى ضبابية غير خطية عرضت لأول مرة، الصيغة التي تحت الدراسة من الأنواع المهمة من معادلات فولتيرا غير الخطية والتي تحوي على أدنى وأعلى دالة. الفضاء المترى الضبابي لهاوس

دورف المعرف على الاعداد الضبابية والفضاءات المترية الضبابي على الدوال المستمرة الضبابية وأيضاً عرفت على الدوال المستمرة الضبابية المطلقة ذات المعيار المنتظم أعطيت بشكل مفصل. كذلك النتائج الظاهرة حققت على الحالات المعلمية لصيغة المسالة الضبابية، بعض الشروط الضرورية والكافية للتقيد ولبنز لمختلف الدوال ومشتقاتها وكذلك استخدام قاعدة لبنز لصيغة المسالة المعروضة. الوجودية والوحدانية المحلية للحلول على فضاء الدوال المستمرة الضبابية المطلقة ذات المعيار المنتظم والذي هو تعميم لفضاء الاعداد الضبابية والمبنية على شرط الانكماش والذي زود من خلال استخدام خواص الفضاء المترية الفضاء المترية المختار. كذلك استخدام مفهوم تكامل ريمان الضبابي ودالة النورم لبرهان نتائج مهمة ووضحت ببعض من الأمثلة لإيضاح المعادلات التكاملية الضبابية غير الخطية المقترحة وحلها في الفضاء المختار والفضاء النورمي وخواصه ويحقق كل الفرضيات للمركبات صيغة المسالة المقترحة وإظهار المقدرات للشروط المناسبة في الفضاء المترية الضبابي المختار الي يعتبر شرط كافي لضمان وجود الحل الضبابي للمعادلات المقترضة.

## 1. Introduction

The fuzzy differential equations are considered as one of the basic equations in the field of engineering modeling, medical and physical applications, as most of their models are defined in fuzzy sets, and therefore the solutions are fuzzy, so they are confined between an upper value and a lower value, and these equations may consist of Volterra and Fredholm integral formulations. Sometime these fuzzy integral equations need some initial conditions or boundary conditions.

The interesting issue in analytic of fuzzy integral equations is the constriction of boundedness as well as necessary and sufficient conditions of convergent, so we presented the following researchers such as some of them studied the sufficient conditions for the solution to be of fuzzy integral equations is bounded in [1]. The approximate solution of nonlinear Hammerstein fuzzy integral equations that is nonlinear, moreover the convergent of successive approximations method have been presented in [2]. The second kind of linear Fredholm fuzzy integral equations instructed their analytic solution by using parametric form of fuzzy numbers, see [3]. In [4], the fuzzy Volterra integral equations with time lag, also the efficient iterative and equivalent of uniqueness and convergence of the method had been given in details. Sometime studied together analytic and approximate solution and presented reliable numerical algorithm. The convert of linear fuzzy Fredholm integral equation to linear system of second kind integral equation is studied with fuzzy number parametric form, [5]. as well as the new interesting results of common fixed-point used in applications on stochastic volterra integral equations achieved contraction prosperity, [6]. In [7], the ordinary approximation technique is used to solve fuzzy Fredholm integral equations with respect fuzzy valued functions is introduced by classical approximation.

The hybrid method for computation solutions of equations with degenerate kernel devoted in [8]. The optimal values of solution appeared by using homotopy analytic method in [9]. The existence and uniqueness are interested in many articles which is established the analytically and approximate solutions in different approaches for different methods where their targets are integrals have types Volterra and Fredholm. [10], [11],[12],

The fuzzy differentiability and integrability studied with a new notion correlated fuzzy processes on time scales was happened in [13], fuzzy Volterra and Fredholm integral equations with the convolution type kernel forced the integral fuzzy problem under Hukuhara differentiability used fuzzy Laplace transform method , [14]. In [6], the mixed of two technical of methods such as fuzzy Lagrange interpolation and the fuzzy Gauss-Legendre quadrature

formula for establish collocation method based are interesting to solve linear and nonlinear fuzzy integral equations of two-dimensional and some main results for convergent

There are many numerical methods such as embedding method, Adomian method, Adomian decomposition method, variational iteration method and homotopy analysis method was used to compute a numerical solution for fuzzy differential and integral equations which investigated with different types of kernels functions and some of them used weakly singular kernels where in the domain of integration may change sign, or some time approximated the fuzzy delay integral equations by piecewise fuzzy polynomial interpolation see [15], [16]. Also the numerical algorithm for solving Fredholm and Volterra Riemann integral second kind and two-dimensional nonlinear fuzzy integrals, all have been presented to convert to a nonlinear system in a crisp case, see [17], [18], [19] and analytic of solution for second kind fuzzy Fredholm integral equations, [20]. The optimal homotopy asymptotic method which is efficient algorithm for solution of system of fuzzy Fredholm integral equations with two dimensional, [21].

The existence and uniqueness of solution for the Volterra-Fredholm nonlinear integral equation with fuzzy parameters have been presented in [22]. In [23], Sharif et al. solved the one-dimensional fuzzy integral equations studded with fuzzy Laplace transform method. Double fuzzy sumudu transform used for solving Volterra fuzzy integral equations were carried by Alidema and Georgieva in [24]. The existence and uniqueness of nonlinear Volterra Fuzzy Integral Equations solutions have been investigated in [11]. Fuzzy integral equation with nonlinear fuzzy kernels has been studied with analysis of convergence which introduced in [25]. In [26], fuzzy integral equations of product type have been studied their existence and uniqueness solution. Additionally, in [8], the Integro-differential equations are nonlinear fuzzy. An Application for fuzzy Integral Equations by new Contraction studied in [12].

The aim of this paper to study the existence and uniqueness solutions of some fuzzy integral equations in uniform modulus of absolutely continuous functions space and explained in details by some illustrative examples simulated the formulation of fuzzy integral equations.

$$\tilde{u}(t, \alpha) = \tilde{f}(t, \alpha) + {}_F(FR) \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \quad (1)$$

where  $\tilde{F}: [a, b] \times [a, b] \times \mathbb{R}_F \times \mathbb{R}_F \rightarrow \mathbb{R}_F$ ,  $\tilde{f}(t, \alpha): [a, b] \times [0, 1] \rightarrow \mathbb{R}_F$ ,  $\tilde{F}_1, \tilde{F}_2: [a, b] \times [a, b] \times \mathbb{R}_F \rightarrow \mathbb{R}_F$  and  $\tilde{u}(s, \alpha): [a, b] \times [0, 1] \rightarrow \mathbb{R}_F$ .

This paper contains four sections: In section 2, some basic preliminaries, namely definitions such as Hausdorff metric on fuzzy number set, the fuzzy space of continuous functions and the first study in this paper is the space of all uniform modulus of absolutely fuzzy continuous as well as the fuzzy matrices defined on these spaces to satisfy the purpose of the Remain fuzzy integrals. In section 3, studied and presented necessary and sufficient hypotheses to achieve existence and uniqueness for classes of fuzzy integral equations with suitable matrices and norms defined on related fuzzy differentials and integrals. In section 4. We give an illustrative example to prove the conditions proposed for existence and uniqueness of some fuzzy integral equations. Lastly, conclusions are shown in section 5.

## 2. Preliminaries

In this section interesting definitions and results used as a basic concept of existence and uniqueness for presented fuzzy integral equations.

**Definition 1, [26].**

If the following properties are satisfied for the function  $u: \mathbb{R} \rightarrow [0,1]$ ,  $\mathbb{R}$  is a real number such that

- (1) there exist  $a_0 \in \mathbb{R}$  such that  $u(a_0) = 1$ . Then  $u$  is normal.
- (2)  $u(\hat{\lambda}a + (1 - \hat{\lambda})b) \geq \min \{u(a), u(b)\}$ ,  $\forall a, b \in \mathbb{R}, 0 \leq \hat{\lambda} \leq 1$ . Then  $u$  is convex fuzzy set.
- (3)  $u$  is the upper semi-continuous on  $\mathbb{R}$ .
- (4) The support  $[u]_0 = \text{cl}\{a \in \mathbb{R}: u(a) > 0\}$  is a compact set.

Then  $u$  is a fuzzy number and the set of all fuzzy numbers is denoted by  $R_F$ .

**Remark 1 [27].**

1. Any real number  $a \in \mathbb{R}$  which can be described as  $\tilde{a} = \chi_{\{a\}}$  and therefore  $\mathbb{R} \subset R_F$ .
2. The  $r$ -level set  $[u]_r = \{a \in \mathbb{R}: u(a) \geq r\}$  that is a closed interval  $[u]_r = [\underline{u}_r, \bar{u}_r]$  for all  $0 \leq r \leq 1$ .
3.  $(\underline{u}(r), \bar{u}(r))$  is a fuzzy number defined as the following  $\underline{u}(r)$  is increasing and  $\bar{u}(r)$  is decreasing functions  $\underline{u}, \bar{u}: [0,1] \rightarrow \mathbb{R}$ .

**Definition 2 [27].**

For  $u, v \in R_F$  and  $\lambda \in \mathbb{R}$ , the sum  $u +_F v$  defined by  $[u + v]_r = [u]_r + [v]_r$  which represents the addition of two intervals (as sub sets of  $\mathbb{R}$ ) and the product  $\lambda u$  defined by  $[\lambda u]_r = \lambda [u]_r$ , for all  $0 \leq r \leq 1$ .

**Definition 3 [28].**

The algebraic properties on  $R_F$  defined as follows:

for any  $u, v, w \in R_F$ , we have that

1.  $u +_F (v +_F w) = (u +_F v) +_F w$  and  $u +_F v = v +_F u$
2.  $u +_F \tilde{0} = \tilde{0} +_F u$  for  $\tilde{0} = \chi_{\{0\}}$
3.  $(a +_F b) \cdot_F u = a \cdot_F u +_F b \cdot_F u$ , for all  $a, b \in \mathbb{R}$  with  $ab \geq 0$ .
4.  $a \cdot_F (u +_F v) = a \cdot_F u +_F a \cdot_F v$  for all  $a \in \mathbb{R}$
5.  $a \cdot_F (b \cdot_F v) = (a \cdot_F b) \cdot_F v$ , for all  $a, b \in \mathbb{R}$  and  $1 \cdot_F u = u$ .

**Definition 4 [29].**

The distance between fuzzy numbers defined by:

$D(u, v) = \sup_{r \in [0,1]} \{\max(|\underline{u}_r - \underline{v}_r|, |\bar{u}_r - \bar{v}_r|)\}$  for all  $u, v \in R_F$  which is called Hausdorff metric.

**Lemma 1 [29].**

The Hausdorff metric has the following properties for all  $u, v, w, e \in R_F$

1.  $(R_F, D)$  is a complete metric space
2.  $D(u +_F w, v +_F w) = D(u, v)$
3.  $D(u +_F v, w +_F e) \leq D(u, w) +_F D(v, e)$
4.  $D(k \cdot_F u, k \cdot_F v) = |k| D(u, v)$ , for all  $k \in \mathbb{R}$ .
5.  $(k_1 \cdot_F u, k_2 \cdot_F u) = |k_1 - k_2| D(u, \tilde{0})$ , for all  $k_1, k_2 \in \mathbb{R}$  with  $k_1 k_2 \geq 0$ .

The property (4) gives the usual norm which is defined by the function  $\|\cdot\|: R_F \rightarrow \mathbb{R}$  such that  $\|u\| = D(u, \tilde{0})$  and we have to use it in throughout the article.

**Lemma 2 [30].**

If  $\hat{f}, \hat{g}: [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}_F$  are fuzzy functions which is continuous, then the function  $F: [a, b] \rightarrow \mathbb{R}^+$  defined by  $F(t) = D(\hat{f}(t), \hat{g}(t))$  is continuous on  $[a, b]$ . Also  $D(\hat{f}(t), \tilde{0}) = \|\hat{f}(t)\| \leq M$  for all  $x \in [a, b]$ ,  $M > 0$  that is  $\hat{f}$  is Globally bounded.

**Definition 5 [30].**

Let  $\tilde{\vartheta}: [a, b] \rightarrow [0, 1]$  be a real valued function, for every partition  $p=[y_0, y_1, \dots, y_m]$  of  $[a, b]$  and every  $\eta_j \in [y_{j-1}, y_j]$ ,  $1 \leq j \leq m$ , we define  $s_q = \sum_{i=0}^m \tilde{\vartheta}(\eta_j)(y_i - y_{i-1})$   
 Now let  $\Delta_j = \max|y_i - y_{i-1}|$ ,  $i=1, 2, \dots, m$  then for  $\tilde{\vartheta}(y)$ , over  $[a, b]$  the defined integral can be written as  $\lim_{\Delta_j \rightarrow 0} s_q = \int_a^b \tilde{\vartheta}(y) dy$  and the parametric form may be represented as  $\int_a^b \tilde{\vartheta}(y) dy = (\int_a^b \underline{\vartheta}(y, k) dy, \int_a^b \bar{\vartheta}(y, k) dy)$

**Lemma 3 [30].**

If (FR) Remain integrable fuzzy functions, and real numbers defined by  $\hat{f}, \hat{g}: [a, b] \rightarrow \mathbb{R}_F$ , then

$$(FR) \int_a^b (\alpha_{\cdot F} \hat{f}(t) + \beta_{\cdot F} \hat{g}(t)) dt = \alpha_{\cdot F} (FR) \int_a^b \hat{f}(t) dt + \beta_{\cdot F} (FR) \int_a^b \hat{g}(t) dt$$

The following definition included the different fuzzy metrics on different fuzzy spaces related among them are interesting to prove the existence of solution for proposal fuzzy nonlinear equation (1), and also achieved interested condition is a completeness.

**Definition 6.**

1. The fuzzy space of continuous functions over  $[a, b]$  denoted by  $C_F([a, b])$ . The metric defined on  $C_F([a, b])$  is  $D^*(f, g) = \sup_{a \leq t \leq b} D(f(t), g(t))$ , for all  $f, g \in C_F([a, b])$  and complete fuzzy metric space. [13]
2. The function  $w_{[a, b]}(f, \delta): \mathbb{R}^+ \cup \{0\} \rightarrow \mathbb{R}^+$  defined by  $w_{[a, b]}(f, \delta) = \sup\{\sum_{i=1}^n D(f(t_i), f(\acute{t}_i)); t_i, \acute{t}_i \in [a, b], \sum_{i=1}^n |t_i - \acute{t}_i| \leq \delta\}$  is called uniform modulus of absolutely fuzzy continuous of  $f \in C_F([a, b])$ ,  $a \leq \acute{t}_i \leq t_i \leq b$ .
3. Let  $f \in V_F([a, b])$  be the space of all uniform modulus of absolutely fuzzy continuous of  $f$  on  $[a, b]$  and  $\int_a^t (D(\frac{\partial}{\partial t} f(t), \tilde{0})) dt$  is a fuzzy Riemann integral.
4. The fuzzy metric defined on  $V_F([a, b])$  as follows:  

$$D^\#(\Gamma_1, \Gamma_2) = D^*(\Gamma_1(t), \Gamma_2(t)) + \int_a^t D^*(\frac{\partial}{\partial x} \Gamma_1(x), \frac{\partial}{\partial x} \Gamma_2(x)) dx,$$
 for any  $\Gamma_1, \Gamma_2 \in V_F([a, b])$  and  $V_F([a, b])$  is a complete metric space corresponding to a complete fuzzy metric space  $(C_F([a, b]), D^*)$ .

The main results of this paper is to prove the existence and uniqueness of Fuzzy integral equations in (1) on  $V_F([a, b])$  be the space of all uniform modulus of absolutely fuzzy continuous of  $f$  on  $[a, b]$  by using metric space  $D^\#(., .)$  defined in definition(6) (4).

**3. Main Results**

In this section the fuzzy problem formulation has been presented with Remain fuzzy integral and fuzzy operations and take a class of Volterra fuzzy nonlinear integral formulation which given to study their existence solution.

Consider the following Fuzzy integral nonlinear equation

$$\tilde{u}(t, \alpha) = \tilde{f}(t, \alpha) +_F (FR) \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds$$

Where  $\tilde{F}: [a, b] \times [a, b] \times R_F \times R_F \rightarrow R_F$ ,  $\tilde{f}(t, \alpha): [a, b] \times [0, 1] \rightarrow R_F$ ,  $\tilde{F}_1, \tilde{F}_2: [a, b] \times [a, b] \times R_F \rightarrow R_F$  and  $\tilde{u}(s, \alpha): [a, b] \times [0, 1] \rightarrow R_F$ .

And their parameter cases are

$$\begin{cases} \underline{u}(t, \alpha) = \underline{f}(t, \alpha) + (FR) \int_a^t F(t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha))) ds \\ \overline{u}(x, \alpha) = \overline{f}(x, \alpha) + (FR) \int_a^t F(t, s, \overline{F}_1(t, s, u(s, \alpha)), \overline{F}_2(t, s, u(s, \alpha))) ds \end{cases} \quad (2)$$

$$\tilde{f}(t, \alpha) = (\underline{f}(t, \alpha), \overline{f}(t, \alpha)), \tilde{u}(s, \alpha) = (\underline{u}(s, \alpha), \overline{u}(s, \alpha)).$$

and

$$\begin{aligned} \overline{F}_1(t, s, u(s, \alpha)) &= \begin{cases} G_1(t, s) \overline{u}(s, \alpha), & G_1(t, s) \geq 0 \\ G_1(t, s) \underline{u}(s, \alpha), & G_1(t, s) < 0 \end{cases} \\ \underline{F}_2(t, s, u(s, \alpha)) &= \begin{cases} G_2(t, s) \overline{u}(s, \alpha), & G_2(t, s) \geq 0 \\ G_2(t, s) \underline{u}(s, \alpha), & G_2(t, s) < 0 \end{cases} \\ \underline{F}_1(t, s, u(s, \alpha)) &= \begin{cases} G_1(t, s) \underline{u}(s, \alpha), & G_1(t, s) \geq 0 \\ G_1(t, s) \overline{u}(s, \alpha), & G_1(t, s) < 0 \end{cases} \\ \overline{F}_2(t, s, u(s, \alpha)) &= \begin{cases} G_2(t, s) \underline{u}(s, \alpha), & G_2(t, s) \geq 0 \\ G_2(t, s) \overline{u}(s, \alpha), & G_2(t, s) < 0 \end{cases} \\ F(t, s, \overline{F}_1(t, s, u(s, \alpha)), \overline{F}_2(t, s, u(s, \alpha))) &= \\ \begin{cases} G_3(t, s) \overline{F}_1(t, s, u(s, \alpha)) \overline{F}_2(t, s, u(s, \alpha)), & G_3(t, s) \geq 0 \\ G_3(t, s) \underline{F}_1(t, s, u(s, \alpha)) \underline{F}_2(t, s, u(s, \alpha)), & G_3(t, s) < 0 \end{cases} \\ F(t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha))) &= \\ \begin{cases} G_3(t, s) \underline{F}_1(t, s, u(s, \alpha)) \underline{F}_2(t, s, u(s, \alpha)), & G_3(t, s) \geq 0 \\ G_3(t, s) \overline{F}_1(t, s, u(s, \alpha)) \overline{F}_2(t, s, u(s, \alpha)), & G_3(t, s) < 0 \end{cases} \end{aligned}$$

$$T(t) = \int_a^t F(t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha))) ds$$

$$L(t) = \int_a^t F(t, s, \overline{F}_1(t, s, u(s, \alpha)), \overline{F}_2(t, s, u(s, \alpha))) ds$$

Let  $T(\cdot) \in C_F([a, b])$ , such that

$$T(t) = (FR) \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds$$

Consider the following hypothesis:

**Hypotheses 1.**

1.  $|G_3(t, s)| \leq M_3$
2.  $\left| \frac{\partial}{\partial t} G_3(t, s) \right| \leq M$
3.  $\left| \frac{\partial}{\partial t} G_1(t, s) \right| \leq M_4$
4.  $\left| \frac{\partial}{\partial t} G_2(t, s) \right| \leq M_5$
5.  $\|\tilde{F}_1(t, s, u(s, \alpha))\| \leq M_1 \|\tilde{u}(s, \alpha)\|$
6.  $\|\tilde{F}_2(t, s, u(s, \alpha))\| \leq M_2 \|\tilde{u}(s, \alpha)\|$
7.  $B(\tilde{u}(t, \alpha)) = \{ \tilde{u}(s, \alpha) : \|\tilde{u}(s, \alpha)\| \leq \hat{r} \}$
8.  $D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) \leq L_1 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
9.  $(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) \leq L_2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
10.  $(\tilde{F}(t, s, \tilde{F}_1(h, s, u_1(s, \alpha)), \tilde{F}_2(h, s, u_1(s, \alpha))), \tilde{F}(t, s, \tilde{F}_1(h, s, u_2(s, \alpha)), \tilde{F}_2(h, s, u_2(s, \alpha))))$

$$\leq L_0 |G_3(t, s)| \left( D \left( \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha)) \right) + D \left( \tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right)$$

11.  $\|G_3(t_i, s) - G_3(\hat{t}_i, s)\| \leq \hat{L}_5 |t_i - \hat{t}_i|$

**Lemma 4.**

Consider the fuzzy integral equation holds the hypotheses (1) as follows:

$$\tilde{u}(t, \alpha) = \tilde{f}(t, \alpha) + \int_a^t F \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds$$

with parameter cases are

$$\begin{cases} \underline{u}(t, \alpha) = \underline{f}(t, \alpha) + \int_a^t F \left( t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha)) \right) ds \\ \bar{u}(x, \alpha) = \bar{f}(x, \alpha) + \int_a^t F \left( t, s, \bar{F}_1(t, s, u(s, \alpha)), \bar{F}_2(t, s, u(s, \alpha)) \right) ds \end{cases}$$

Then the following statements are satisfied:

1.  $D \left( \tilde{F}(t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha))) \right) \leq M_3 M_1 M_2 \hat{r}^2$
2.  $D \left( \tilde{F}(t, s, \tilde{F}_1(h, s, u_1(s, \alpha)), \tilde{F}_2(h, s, u_1(s, \alpha))), \tilde{F}(t, s, \tilde{F}_1(h, s, u_2(s, \alpha)), \tilde{F}_2(h, s, u_2(s, \alpha))) \right) \leq L_0 M_3 (L_1 + L_2) D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
3.  $D \left( \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha))), \tilde{0} \right) \leq \hat{r}^2 (M M_1 M_2 + M_3 (M_4 M_2 + M_5 M_1))$
4.  $D \left( \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))) - \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_2(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))), \tilde{0} \right) \leq (M M_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + M M_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1) D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$

**Proof**

1.  $D \left( \tilde{F}(t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha))), \tilde{0} \right) \leq |G_3(t, s)| \| \tilde{F}_1(t, s, u(s, \alpha)) \| \| \tilde{F}_2(t, s, u(s, \alpha)) \| \leq M_3 M_1 M_2 \hat{r}^2$
2.  $D \left( \tilde{F}(t, s, \tilde{F}_1(h, s, u_1(s, \alpha)), \tilde{F}_2(h, s, u_1(s, \alpha))), \tilde{F}(t, s, \tilde{F}_1(h, s, u_2(s, \alpha)), \tilde{F}_2(h, s, u_2(s, \alpha))), \tilde{0} \right) \leq L_0 |G_3(t, s)| \left( D \left( \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha)) \right) + D \left( \tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right) \leq L_0 M_3 (L_1 + L_2) D(u_1(s, \alpha), u_2(s, \alpha))$
3.  $D \left( \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha))), \tilde{0} \right) = \left\| \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha))) \right\| \leq \left| \frac{\partial}{\partial t} G_3(t, s) \right| \| \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(h, s, u(s, \alpha)) \| + |G_3(t, s)| \left\| \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(h, s, u(s, \alpha)) \right) \right\| \leq M \| \tilde{F}_1(h, s, u(s, \alpha)) \tilde{F}_2(h, s, u(s, \alpha)) \| + |G_3(t, s)| \left\| \frac{\partial}{\partial t} G_1(t, s) u(s, \alpha) \tilde{F}_2(h, s, u(s, \alpha)) + \tilde{F}_1(h, s, u(s, \alpha)) \frac{\partial}{\partial t} G_2(t, s) u(s, \alpha) \right\| \leq (M M_1 M_2 + M_3 (M_4 M_2 + M_5 M_1)) \| \tilde{u}(s, \alpha) \|^2 \leq \hat{r}^2 (M M_1 M_2 + M_3 (M_4 M_2 + M_5 M_1))$   
 for  $t, s \in [a, b]$  and  $M_k > 0$ .

4.

$$\begin{aligned}
 & D \left( \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))) (-1)_{\cdot F} \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_2(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))), \tilde{0} \right) = \\
 & D \left( \frac{\partial}{\partial t} G_3(t, s) \tilde{F}_1(t, s, u_1(s, \alpha)) \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \frac{\partial}{\partial t} G_3(t, s) \tilde{F}_1(t, s, u_1(s, \alpha)) \right. \\
 & \quad \left. \tilde{F}_2(t, s, u_2(s, \alpha)) + {}_F \frac{\partial}{\partial t} G_3(t, s) \tilde{F}_1(t, s, u_1(s, \alpha)) \tilde{F}_2(t, s, u_2(s, \alpha)) - \right. \\
 & \quad \left. \frac{\partial}{\partial t} G_3(t, s) \tilde{F}_1(t, s, u_2(s, \alpha)) \tilde{F}_2(t, s, u_2(s, \alpha)), \tilde{0} \right) \\
 & = D \left( \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right. \\
 & \quad \left. + {}_F \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) - \tilde{F}_1(t, s, u_2(s, \alpha)) \right), \tilde{0} \right) = \\
 & D \left( \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right. \\
 & \quad \left. + {}_F G_3(t, s) \frac{\partial}{\partial t} \left( \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) - \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right) \right) = \\
 & D \left( \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) - \tilde{F}_2(t, s, u_2(s, \alpha)) \right) + {}_F \right. \\
 & \quad \left. G_3(t, s) \left( \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right) \right. \\
 & \quad \left. + {}_F G_3(t, s) \left( \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) - \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right) \right) \\
 & \quad \left. + {}_F G_3(t, s) \left( \left( \tilde{F}_2(t, s, u_1(s, \alpha)) \right) \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right) \right) \\
 & \quad \left. \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) + {}_F \right. \\
 & \quad \left. G_3(t, s) \left( \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right) \right), \tilde{0} = \\
 & \left\| \frac{\partial}{\partial t} G_3(t, s) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right\| \\
 & \quad + {}_F G_3(t, s) \left\| \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right\| \\
 & \quad + {}_F G_3(t, s) \left\| \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right\| \\
 & \quad + {}_F \frac{\partial}{\partial t} + {}_F G_3(t, s) \left\| \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) \right) \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right\| \\
 & \quad + {}_F G_3(t, s) \left\| \left( \tilde{F}_2(t, s, u_1(s, \alpha)) \right) \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right\| \\
 & \leq \left\| \frac{\partial}{\partial t} G_3(t, s) \right\| \left\| \tilde{F}_1(t, s, u_1(s, \alpha)) \right\| \left\| \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right\| + \\
 & \left\| G_3(t, s) \right\| \left\| \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) \right) \right\| \left\| \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right\| + \\
 & \left\| G_3(t, s) \right\| \left\| \tilde{F}_1(t, s, u_1(s, \alpha)) \right\| \left\| \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_2(t, s, u_2(s, \alpha)) \right) \right\| + \\
 & \left\| \frac{\partial}{\partial t} G_3(t, s) \right\| \left\| \tilde{F}_2(t, s, u_1(s, \alpha)) \right\| \left\| \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right\| + \\
 & \left\| G_3(t, s) \right\| \left\| \frac{\partial}{\partial t} \left( \tilde{F}_2(t, s, u_1(s, \alpha)) \right) \right\| \left\| \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right\| + \\
 & \left\| G_3(t, s) \right\| \left\| \tilde{F}_2(t, s, u_1(s, \alpha)) \right\| \left\| \frac{\partial}{\partial t} \left( \tilde{F}_1(t, s, u_1(s, \alpha)) (-1)_{\cdot F} \tilde{F}_1(t, s, u_2(s, \alpha)) \right) \right\| \leq \\
 & MM_1 \hat{r} L_2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) + \\
 & M_3 M_5 \hat{r} L_2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) + M_3 M_1 \hat{r} L_2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) \\
 & + MM_2 \hat{r} L_1 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) + M_3 M_5 \hat{r} L_1 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) + \\
 & M_3 M_2 \hat{r} M_1 L_1 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) \leq (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + \\
 & M_3 M_5 \hat{r} L_1) D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) \text{ and for any } \tilde{u}_1(t, s, \alpha), \tilde{u}_2(t, s, \alpha) \text{ are fuzzy valued continuous} \\
 & \text{functions.}
 \end{aligned}$$



**Theorem 1.**

If  $\hat{T}(t) \in C_F([a, b])$  such that

$$\hat{T}(t) = (RF) \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(h, s, u(s, \alpha)), \tilde{F}_2(h, s, u(s, \alpha)) \right) ds.$$

Then  $\hat{T}(t)$  is a uniform modulus of absolutely continuous on  $[a, b]$ .

**Proof**

$$\begin{aligned} & \sum_{i=1}^n D(T(t_i), T(\hat{t}_i), \tilde{0}) \\ = & \sum_{i=1}^n D \left( (RF) \int_a^{t_i} \tilde{F}(t, s, \tilde{F}_1(h, s, u(s, \alpha)), \tilde{F}_2(h, s, u(s, \alpha))) ds \right. \\ & \left. (RF) \int_a^{\hat{t}_i} \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds, \tilde{0} \right) \\ \text{By lemma (2.7), we get that} \\ & \sum_{i=1}^n \left\| (RF) \int_a^{t_i} \tilde{F} \left( t_i, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds (-1)_{\cdot F} \right. \\ & \left. (RF) \int_a^{\hat{t}_i} \tilde{F} \left( \hat{t}_i, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \right\| \\ = & \sum_{i=1}^n \left\| (RF) \int_a^{t_i} \tilde{F} \left( t_i, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \right. \\ & \left. (-1)_{\cdot F} (RF) \int_a^{\hat{t}_i} \tilde{F} \left( \hat{t}_i, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \right\| \\ & \sum_{i=1}^n \left\| (RF) \int_a^{t_i} G_3(t_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right. \\ & \left. (-1)_{\cdot F} (RF) \int_a^{\hat{t}_i} G_3(\hat{t}_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right\| \\ \leq & \sum_{i=1}^n \left\| (RF) \int_a^{t_i} G_3(t_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right. \\ & \left. +_F (RF) \int_{\hat{t}_i}^{t_i} G_3(t_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right. \\ & \left. (-1)_{\cdot F} (RF) \int_a^{\hat{t}_i} G_3(\hat{t}_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right\| \\ \leq & \sum_{i=1}^n \left\| (RF) \int_a^{t_i} G_3(t_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right. \\ & \left. (-1)_{\cdot F} (RF) \int_a^{\hat{t}_i} G_3(\hat{t}_i, s) \tilde{F}_1(t, s, u(s, \alpha)) \tilde{F}_2(t, s, u(s, \alpha)) ds \right\| \\ & + \sum_{i=1}^n \left\| (RF) \int_{\hat{t}_i}^{t_i} G_3(t_i, s) \tilde{F}_1(h, s, u(s, \alpha)) \tilde{F}_2(h, s, u(s, \alpha)) ds \right\| \\ \leq & \sum_{i=1}^n \int_a^{t_i} \|G_3(t_i, s) - G_3(\hat{t}_i, s)\| \| \tilde{F}_1(t, s, u(s, \alpha)) \| \| \tilde{F}_2(t, s, u(s, \alpha)) \| ds + \sum_{i=1}^n \int_{\hat{t}_i}^{t_i} \|G_3(t_i, s) - \\ & G_3(\hat{t}_i, s)\| \| \tilde{F}_1(t, s, u(s, \alpha)) \| \| \tilde{F}_2(t, s, u(s, \alpha)) \| ds \\ \leq & \hat{L}_5 \sum_{i=1}^n |t_i - a| |t_i - \hat{t}_i| M_1 M_2 \| \tilde{u}(s, \alpha) \|^2 + \hat{L}_5 \sum_{i=1}^n |t_i - \hat{t}_i|^2 M_1 M_2 \| \tilde{u}(s, \alpha) \|^2 \leq \\ & \hat{L}_5 \sum_{i=1}^n |b - a| |t_i - \hat{t}_i| M_1 M_2 \tilde{r}^2 \\ \leq & 2 \hat{L}_5 |b - a|^2 M_1 M_2 \tilde{r}^2 \left( \frac{\epsilon}{2 \hat{L}_5 |b - a|^2 M_1 M_2 \tilde{r}^2} \right) \leq \epsilon \end{aligned}$$

Hence  $\hat{T}(t)$  a uniform modulus of absolutely continuous on  $[a, b]$ .

**Theorem 2.**

Let  $\hat{T} \in C_F([a, b])$  such that

$$\frac{\partial}{\partial t} \hat{T}(t) = \frac{\partial}{\partial t} \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \tag{3}$$

Then  $\int_a^b \left( D \left( \frac{\partial}{\partial t} \hat{T}(t), \tilde{0} \right) \right) dt$  is fuzzy Riemann integrable .

**Proof**

We have that  $\frac{\partial}{\partial t} \hat{T}(t) = \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) + \int_a^t \frac{\partial}{\partial t} \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds$  then for a fuzzy number  $\tilde{A}$ , we have that

$$\begin{aligned}
 & D \left( \int_a^t \left( D \left( \frac{\partial}{\partial t} \hat{T}(t), \tilde{0} \right) \right) dt, \tilde{A} \right) = \\
 & D \left( \int_a^t \left\| \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) + {}_F \int_a^t \frac{\partial}{\partial t} \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds \right\| dt, \tilde{A} \right) \\
 & \leq D \left( \int_a^t \left\| \tilde{F} \left( t, s, \tilde{F}_1(h, s, u(s, \alpha)), \tilde{F}_2(h, s, u(s, \alpha)) \right) \right\| dt, \tilde{A} \right) \\
 & \quad + D \left( \int_a^t \left\| \int_a^t \frac{\partial}{\partial t} \tilde{F} \left( t, s, \tilde{F}_1(h, s, u(s, \alpha)), \tilde{F}_2(h, s, u(s, \alpha)) \right) ds \right\| dt, \tilde{A} \right)
 \end{aligned}$$

Where  $\tilde{A}$  is a fuzzy Number .

By condition (1) and (2) , we get

$$\begin{aligned}
 & D \left( \int_a^t \left( D \left( \frac{\partial}{\partial t} \hat{T}(t), \tilde{0} \right) \right) dt, A \right) \leq M_3 M_1 M_2 \hat{r}^2 \int_a^b dt + \\
 & \int_a^t \left\| \int_a^t \frac{\partial}{\partial t} \tilde{F} \left( t, s, \tilde{F}_1(h, s, u(s, \alpha)), \tilde{F}_2(h, s, u(s, \alpha)) \right) ds \right\| dt
 \end{aligned}$$

Hence,

$$\int_a^t \left( D \left( \frac{\partial}{\partial t} T(t), \tilde{0} \right) \right) dt \leq M_3 M_1 M_2 \hat{r}^2 (b - a) + (b - a) \hat{r}^2 (M M_1 M_2 + M_3 (M_4 M_2 + M_5 M_1)) < \infty$$

Hence,  $\int_a^t \left( D \left( \frac{\partial}{\partial t} \hat{T}(t), \tilde{0} \right) \right) dt$  is fuzzy Riemann integrable.

Let a sub-intervals  $a \leq x_0 \leq x_1 \dots \leq x_n \leq b$  of  $[a, b]$  into  $N$ , such that  $\Delta x = x_j - x_{j-1}, j = 1, \dots, N$ , and  $\Delta x = \frac{b-a}{N}$ , this partition are needed in following result.

**Theorem 3.**

Let  $\Gamma \in V_F([x_j, x_{j+\Delta x}])$  such that

$$\Gamma(\tilde{u}_1(s, \alpha)) = \tilde{f}(t, \alpha) + {}_F(FR) \int_a^t \tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))) ds$$

$$\Gamma(\tilde{u}_2(s, \alpha)) = \tilde{f}(t, \alpha) + {}_F(FR) \int_a^t \tilde{F}(t, s, \tilde{F}_1(t, s, u_2(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) ds$$

Then  $D^\#(\Gamma(\tilde{u}_1), \Gamma(\tilde{u}_2)) \leq D^*(\tilde{u}_1, \tilde{u}_2)$  for all  $\tilde{u}_1, \tilde{u}_2 \in C_F([a, b])$ , and  $\alpha \in [0,1]$ , and  $\Gamma$  has locally existence and unique solution.

**Proof**

We have that

$$\begin{aligned}
 & D^\#(\Gamma(\tilde{u}_1), \Gamma(\tilde{u}_2)) = D^*(\Gamma_1(t, \alpha), \Gamma_2(t, \alpha)) + {}_F \int_{t_j}^{t_j+\Delta t} D^* \left( \frac{\partial}{\partial t} \Gamma_1(t, \alpha), \Gamma_2(t, \alpha) \right) dt \\
 & = \sup_{a \leq t \leq b} D(\Gamma(\tilde{u}_1(s, \alpha))(t), \Gamma(\tilde{u}_2(s, \alpha))(t)) \\
 & + {}_F(FR) \int_{t_j}^{t_j+\Delta t} \sup_{a \leq t \leq b} D \left( \frac{\partial}{\partial t} \Gamma(\tilde{u}_1(s, \alpha))(t), \frac{\partial}{\partial t} \Gamma(\tilde{u}_2(s, \alpha))(t) \right) dt \\
 & = (FR) \int_{t_j}^{t_j+\Delta t} \sup_{a \leq t \leq b} D \left( \frac{\partial}{\partial t} \Gamma(\tilde{u}_1(s, \alpha))(t), \Gamma(\tilde{u}_2(s, \alpha))(t) \right) dt \\
 & = (FR) \int_{t_j}^{t_j+\Delta t} \left( \sup_{a \leq t \leq b} D(\tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))) \right. \\
 & \quad , \tilde{F}(t, s, \tilde{F}_1(t, s, u_2(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) \left. \right) ds + {}_F(FR) \sup_{a \leq t \leq b} \\
 & \int_{t_j}^t D \left( \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))) \right. \\
 & \quad , \left. \frac{\partial}{\partial t} \tilde{F}(t, s, \tilde{F}_1(t, s, u_2(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) \right) ds \Big) dt \leq \int_{t_j}^{t_j+\Delta t} \left( L_0 M_3 (L_1 + \right. \\
 & L_2) D^*(\tilde{u}_1(t, \alpha), \tilde{u}_2(t, \alpha)) dt + \int_{t_j}^t (M M_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + M M_2 \hat{r} L_1 + \\
 & M_3 M_5 \hat{r} L_1) D^*(\tilde{u}_1(t, \alpha), \tilde{u}_2(t, \alpha)) ds \Big) dt
 \end{aligned}$$

$$\leq \Delta t \left( L_0 M_3 (L_1 + L_2) + (\Delta t L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)) \right) D^*(\tilde{u}_1(t, \alpha), \tilde{u}_2(t, \alpha))$$

Set  $\Delta t < 1$ ,  $\sigma(t) = \left( L_0 M_3 (L_1 + L_2) + (\Delta t L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)) \right)$  and  $\left( L_0 M_3 (L_1 + L_2) + (\Delta t L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)) \right) < \left( L_0 M_3 (L_1 + L_2) + (L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)) \right)$

Take  $\Delta t < \frac{1}{L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)}$ , therefore

$$D^\#(\Gamma(\tilde{u}_1), \Gamma(\tilde{u}_2)) \leq \Delta t \sigma(t) D^*(\tilde{u}_1(t, \alpha), \tilde{u}_2(t, \alpha)) < \Delta t \left( L_0 M_3 (L_1 + L_2) + (L_3 (MM_1 \hat{r} L_2 + M_3 M_1 \hat{r} \hat{M} + M_3 \hat{M}_1 \hat{r} M_5 L_2 + MM_2 \hat{r} L_1 + M_3 M_5 \hat{r} L_1)) \right) D^*(u_1(t, \alpha), u_2(t, \alpha)) <$$

$D^*(\tilde{u}_1(t, \alpha), \tilde{u}_2(t, \alpha))$ , hence

$$D^\#(\Gamma(\tilde{u}_1), \Gamma(\tilde{u}_2)) \leq D^*(\tilde{u}_1, \tilde{u}_2).$$

Hence  $\Gamma \in V_F([x_j, x_{j+\Delta x}])$  for arbitrary  $\Delta x$  is a contractive mapping and since  $V_F([x_j, x_{j+\Delta x}])$  is a complete fuzzy metric space so by Banach fixed point theorem  $\Gamma$  has locally existence and unique solution in  $V_F([x_j, x_{j+\Delta x}])$ .

The following illustrative examples are presented to show the role of necessary and sufficient conditions to grantee the existence and uniqueness of solution.

#### 4. Illustrative Examples:

##### Example 1.

Consider the following fuzzy integral nonlinear equation

$$\tilde{u}(t, \alpha) = \tilde{f}(t, \alpha) + \int_a^t \tilde{F} \left( t, s, \tilde{F}_1(t, s, u(s, \alpha)), \tilde{F}_2(t, s, u(s, \alpha)) \right) ds$$

Has the following upper and lower parts such as

$$\begin{cases} \underline{u}(t, \alpha) = \underline{f}(t, \alpha) + \int_a^t \underline{F} \left( t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha)) \right) ds \\ \bar{u}(t, \alpha) = \bar{f}(t, \alpha) + \int_a^t \bar{F} \left( t, s, \bar{F}_1(t, s, u(s, \alpha)), \bar{F}_2(t, s, u(s, \alpha)) \right) ds \end{cases}$$

Where

$$\underline{f}(t, \alpha) = \frac{37 \alpha^2 t^{10}}{224},$$

$$\bar{f}(t, \alpha) = \frac{37 \left(\frac{\alpha}{2} - 2\right)^2 t^{10}}{504}, \text{ and } G_3(t, s) = st$$

$$\underline{F}_1(t, s, u(s, \alpha)) = G_1(t, s) \underline{u}(t, \alpha) = (t + s) \underline{u}(t, \alpha)$$

$$\underline{F}_2(t, s, u(s, \alpha)) = G_2(t, s) \underline{u}(t, \alpha) = (t^2 - s^2) \underline{u}(t, \alpha), \text{ also}$$

$$\bar{F} \left( t, s, \bar{F}_1(t, s, u(s, \alpha)), \bar{F}_2(t, s, u(s, \alpha)) \right) = st(t + s) \bar{u}(t, \alpha) (t^2 - s^2) \bar{u}(t, \alpha)$$

and we have the following upper deceptions of functions

$$\bar{F}_1(t, s, u(s, \alpha)) = (t + s) \bar{u}(t, \alpha)$$

$$\bar{F}_2(t, s, u(s, \alpha)) = (t^2 - s^2) \bar{u}(t, \alpha), \text{ also}$$

$$\bar{F} \left( t, s, \bar{F}_1(t, s, u(s, \alpha)), \bar{F}_2(t, s, u(s, \alpha)) \right) = st(t + s) \bar{u}(t, \alpha) (t^2 - s^2) \bar{u}(t, \alpha)$$

Which satisfies the following conditions:

- 1)  $|G_3(t, s)| = |st| \leq b^2 = M_3$
- 2)  $\left| \frac{\partial}{\partial t} G_3(t, s) \right| = |s| \leq b = M$
- 3)  $\left| \frac{\partial}{\partial t} G_1(t, s) \right| \leq 1 = M_4$
- 4)  $\left| \frac{\partial}{\partial t} G_2(t, s) \right| \leq 2b = M_5$
- 5)  $\|\tilde{F}_1(t, s, u(s, \alpha))\| \leq 2b\|\tilde{u}(s, \alpha)\| = M_1\|\tilde{u}(s, \alpha)\|$
- 6)  $\|\tilde{F}_2(t, s, u(s, \alpha))\| \leq b^2\|\tilde{u}(s, \alpha)\| = M_2\|\tilde{u}(s, \alpha)\|$
- 7)  $\mathbf{B}(u(t, \alpha)) = \{u(t, \alpha) : \|u(t, \alpha)\| \leq \hat{r}\}$ ,  $\hat{r}$  is a radius fuzzy number.
- 8)  $D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) \leq 2bD(u_1(s, \alpha), u_2(s, \alpha)) = L_1D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
- 9)  $(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) \leq b^2D(u_1(s, \alpha), u_2(s, \alpha)) = L_2D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
- 10)  $D(\tilde{F}(t, s, \tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_1(s, \alpha))), \tilde{F}(t, s, \tilde{F}_1(h, s, u_2(s, \alpha)), \tilde{F}_2(h, s, u_2(s, \alpha)), 0) \leq 2b^2(D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) + D(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))))$   
 $= L_0|G_3(t, s)| (D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) + D(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))))$
- 11)  $\|G_3(t_i, s) - G_3(\hat{t}_i, s)\| \leq b|t_i - \hat{t}_i| = \hat{L}_5|t_i - \hat{t}_i|$ .

The fuzzy integral equation has a locally existence and unique solution.

### Example 2.

Consider the details components of other fuzzy integral nonlinear equation

$$\underline{f}(x, \alpha) = \frac{41 \alpha^2 t^{12}}{140},$$

$$\bar{f}(x, \alpha) = \frac{41 (\frac{\alpha}{2} - 2)^2 t^{12}}{315}, \text{ and } G_3(t, s) = st^2$$

$$\underline{F}_1(t, s, u(s, \alpha)) = G_1(t, s) \underline{u}(t, \alpha) = (t + s)^2 \underline{u}(t, \alpha)$$

$$\underline{F}_2(t, s, u(s, \alpha)) = G_2(t, s) \underline{u}(t, \alpha) = (t^2 - s^2) \underline{u}(t, \alpha), \text{ also}$$

$$\underline{F}(t, s, \underline{F}_1(t, s, u(s, \alpha)), \underline{F}_2(t, s, u(s, \alpha))) = st^2(t + s)^2 \underline{u}(t, \alpha) (t^2 - s^2) \underline{u}(t, \alpha)$$

and we have the following upper deceptions of functions

$$\overline{F}_1(t, s, u(s, \alpha)) = (t + s)^2 \overline{u}(t, \alpha)$$

$$\overline{F}_2(t, s, u(s, \alpha)) = (t^2 - s^2) \overline{u}(t, \alpha), \text{ also}$$

$$\overline{F}(t, s, \overline{F}_1(t, s, u(s, \alpha)), \overline{F}_2(t, s, u(s, \alpha))) = st^2(t + s)^2 \overline{u}(t, \alpha) (t^2 - s^2) \overline{u}(t, \alpha)$$

Which satisfies the following conditions:

- 1)  $|G_3(t, s)| = |st^2| \leq b^3 = M_3$
- 2)  $\left| \frac{\partial}{\partial t} G_3(t, s) \right| = |2ts| \leq 2b^2 = M$
- 3)  $\left| \frac{\partial}{\partial t} G_1(t, s) \right| \leq 4b = M_4$
- 4)  $\left| \frac{\partial}{\partial t} G_2(t, s) \right| \leq 2b = M_5$
- 5)  $\|\tilde{F}_1(t, s, u(s, \alpha))\| \leq 4b\|\tilde{u}(s, \alpha)\| = M_1\|\tilde{u}(s, \alpha)\|$

- 6)  $\|\tilde{F}_2(t, s, u(s, \alpha))\| \leq b^2 \|\tilde{u}(s, \alpha)\| = M_2 \|\tilde{u}(s, \alpha)\|$
- 7)  $\mathbf{B}(u(t, \alpha)) = \{ \tilde{u}(t, \alpha) : \|\tilde{u}(t, \alpha)\| \leq \hat{r} \}$ ,  $\hat{r}$  is a radius fuzzy number.
- 8)  $D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) \leq b^3 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) = L_1 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
- 9)  $(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))) \leq b^2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha)) = L_2 D(\tilde{u}_1(s, \alpha), \tilde{u}_2(s, \alpha))$
- 10)  $(F(t, s, \tilde{F}_1(h, s, u_1(s, \alpha)), \tilde{F}_2(h, s, u_1(s, \alpha))), \tilde{F}(t, s, \tilde{F}_1(h, s, u_2(s, \alpha)), \tilde{F}_2(h, s, u_2(s, \alpha))), )$   
 $\leq 2b^3 (D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) + D(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))))$   
 $= L_0 |G_3(t, s)| (D(\tilde{F}_1(t, s, u_1(s, \alpha)), \tilde{F}_1(t, s, u_2(s, \alpha))) + D(\tilde{F}_2(t, s, u_1(s, \alpha)), \tilde{F}_2(t, s, u_2(s, \alpha))))$
- 11)  $\|G_3(t_i, s) - G_3(\hat{t}_i, s)\| \leq 2b^2 |t_i - \hat{t}_i| = \hat{L}_5 |t_i - \hat{t}_i|$

Hence the fuzzy integral equation has locally existence and uniqueness solution.

## 5. Conclusion

1. We concluded to establish a fuzzy metric space of uniform modulus of absolutely fuzzy continuous functions need to define a metric space of fuzzy continuous functions and this defined on a set of fuzzy numbers which denoted by  $R_F$ .
2. We concluded to cover all the parameters cases of all components of proposal fuzzy integral problem formulation, we need to explain these classes with upper and lower functions and their ordinary kernel functions which all depended on the values of  $\alpha \in [0,1]$ .
3. We concluded the certain necessary and sufficient conditions was choice it in a good accuracy suitable with the certain hypotheses for proposal problem to investigate the results of existence and uniqueness fuzzy solutions of proposal fuzzy integral classes.
4. We concluded the interesting result is to achieve the metric space of fuzzy continuous functions a fuzzy Riemann integrable which is made a great role for techniques of the proving of main result.
5. We concluded that to be the main results that are interesting for applications in fuzzy theory, we need the illustrative examples and supported that the proposal fuzzy formulation is exist and has a fuzzy solution in suitable fuzzy metric space.
6. We concluded the fuzzy counteractive condition is satisfied by divided of interval solution to grantee the result existence and uniqueness by using fixed point theorem with introduced fuzzy metric space.
7. We concluded that the norm function was defined with respect to the metric space of fuzzy continuous and used in technical of prove some results.
8. We concluded the main result does not satisfy without fixed point theorem on fuzzy numbers that which appearing in all definitions of the proving stricture.

## References

- [1] S. Abbasbandy E. Babolian and M. Alavi, Numerical method for solving linear Fredholm fuzzy integral equations of the second kind, Chaos, Solitons & Fractals, Vol 31, Issue, January 2007, Pages 138-146.
- [2] M. Al smadi, Reliable numerical algorithm for handling fuzzy integral equations of second kind in Hilbert space, Flomat 2019 vol 33, issue 33, pages 583-597.
- [3] A. Mihai, C. Popescu, Approximating the solution of nonlinear Hammerstein fuzzy integral equations, Fuzzy Sets and Systems, Vol 245, 16 June 2014, Pages 1-17.
- [4] T.L. Yookesh, M. Chithambarathanu, E. Boopathi Kumar, P. Sudam Sekhar, Efficiency of iterative filtering method for solving Volterra fuzzy integral equations with a delay and material investigation, Vol 47, Part 17, 2021, Pages 6101-6104.

- [5] J. Nieto, R Rodríguez, Bounded solutions for fuzzy differential and integral equations, *Chaos, Solitons & Fractals*, Vol 27, Issue 5, March 2006, Pages 1376-1386.
- [6] P. Deng, J. Huang, X. Wen, and H. Liu, Discrete collocation method for solving two-dimensional linear and nonlinear fuzzy Volterra integral equations, *Applied Numerical Mathematics*, Vol 171, January 2022, Pages 389-407.
- [7] B. Shiri, I. Perfilieva, and Z. Alijani, Classical approximation for fuzzy Fredholm integral equation *Fuzzy Sets and Systems*, Vole 404, 1 February 2021, Pages 159-177.
- [8] A. Ullah, Z. Ullah. A hybrid method for solving fuzzy Volterra integral equations of separable type kernels, *Journal of King Saud University - Science*, Vol 33, Issue 1, January ,2021, 101-246.
- [9] M. Ali F. Araghi and S. Noeiaghdam, Finding Optimal Results in the Homotopy Analysis Method to Solve Fuzzy Integral Equations, *Advances in fuzzy integral and differential equations*, 2021, pp 173–195.
- [10] S. H. Ghasemi, “Fuzzy Fredholm-Volterra integral equations and existence and uniqueness of solution of them,” *Aust. J. Basic Appl. Sci.*, vol. 5, no. 4, pp. 1–8, 2011.E.
- [11] Hussain and A. W. Ali, “Existence and Uniqueness of The Solution of Nonlinear Volterra Fuzzy Integral Equations,” *Ibn AL-Haitham J. Pure Appl. Sci.*, vol. 26, no. 2, pp. 348–357, 2017.
- [12] V. Samadpour K. Mahaleh and R. Ezzati, “Existence and uniqueness of solution for fuzzy integral equations of product type,” *Soft Comput.*, vol. 25, no. 21, pp. 13287–13295, 2021.
- [13] M. Shahidi, E. Esmi and L.C. Barros, a study on fuzzy Volterra integral equations for correlated fuzzy processes on time scales, *Fuzzy Sets and Systems*, Volume 471, 15,
- [14] A. A. Sharif, M. M. Hamood, and A. D. Khandale, Usage of the Fuzzy Laplace Transform Method for Solving One-Dimensional Fuzzy Integral Equations, November 2023, 108695.
- [15] M. Baghmisheh and R. Ezzati, Application of Hybrid Bernstein Polynomials and Block-Pulse Functions for Solving Nonlinear Fuzzy Fredholm Integral Equations. *Fuzzy Information and Engineering*, 2023. 15(1): 69-86.
- [16] Z. Alijani and U. Kangro, Numerical solution of a linear fuzzy Volterra integral equation of the second kind with weakly singular kernels, *Fuzzy system and mathematics*, Vol 26, pages 12009-12022(2022).
- [17] M. Friedman M. Ming and A. Kandel, Solutions to fuzzy integral equations with arbitrary kernels, *International Journal of Approximate Reasoning* Vol 20, Issue 3, March 1999, Pages 249-262.
- [18] A. Georgieva, Solving two-dimensional nonlinear volterra- Fredholm fuzzy integral equations by adomian decomposition method, *Dynamic Systems and Applications*, 27, No. 4 (2018), 819-835.
- [19] A. Georgieva and D. Gruyter, Solving two-dimensional nonlinear fuzzy Volterra integral equations by homotopy analysis method, *journal Demonstratio Mathematica*, 2021. Vol 54, issue 1.
- [20] A. Molabahrami, A. Shidfar and A. Ghyas, an analytical method for solving linear Fredholm fuzzy integral equations of the second kind, *Computers & Mathematics with Applications*, Vol 61, Issue 9, May 2011, Pages 2754-2761.
- [21] T. Deng, J. Huang, H. Liu and H. Li, Approximate solutions of fuzzy delay integral equations with weakly singular kernels by piecewise fuzzy polynomial interpolation, *Fuzzy*, Vol 470, 30 October 2023, 108652
- [22] S. Q. Hasan and A. J. Abdulqader, “Numerical and analytic method for solving proposal fuzzy nonlinear volterra integral equation by using homotopy analysis method,” *Am. J. Appl. Math.*, vol. 4, no. 3, pp. 142–157, 2016.
- [23] A. A. Sharif, M. M. Hamood, and A. D. Khandagale, “Usage of the Fuzzy Laplace Transform Method for Solving One-Dimensional Fuzzy Integral Equations,” *Equations*, vol. 2, pp. 31–37, 2022.
- [24] A. Alidema and A. Georgieva, “Applications of the double fuzzy Sumudu transform for solving Volterra fuzzy integral equations,” in *AIP Conference Proceedings*, 2019, vol. 2172, no. 1, p. 60006.
- [25] L. Hooshangian, “Analysis of convergence of solution of general fuzzy integral equation with nonlinear fuzzy kernels,” *Theory Approx. Appl.*, vol. 11, no. 2, pp. 97–109, 2017.
- [26] T. Pathinathan., K. Ponnivalavan. and E. Mike Dison., Different Types of Fuzzy Numbers and Certain Properties, *Journal of Computer and Mathematical sciences*, Vol.6(11),631-651, November 2015

- [27] [H. C. Wu, The improper fuzzy Riemann integral and its numerical integration, information science 111(1999)109-137.
- [28] A. Kaufmann and M.M Gupta., Introduction fuzzy arithmetic, Van No strand Reinhold, New York, (1985).
- [29] C. Wu and Z. Gong, On Henstock integral of fuzzy-number functions(I), Fuzzy set syst, vol 120mno. 3, pp 523-532, 2001
- [30] R. Goetschel and W. Vaxman., Elementary fuzzy calculus fuzzy sets and systems,81 (1986),31-43.