



ISSN: 0067-2904

R^* -Small Submodules and R^* -Hollow Modules

Mustafa Mohammed Jasim*, Wasan Khalid

Mathematics Department, College of Science, University of Baghdad, Iraq

Received: 31/10/2023 Accepted: 29/1/2024 Published: xx

Abstract

The purpose of this study is to present a new concepts on a module M over a ring, these concepts is called R^* -small sub module, R^* -hollow module which present generalizations of the small submodule and hollow module, respectively. Key characteristics of these concepts such as the image and direct sum of R^* -small sub module, R^* -hollow module are R^* -small submodule, R^* -hollow submodule, respectively.

Keywords: e^* -essential submodule, R^* -small submodule, R^* -hollow module

المقاسات الجزئية الصغيرة من النمط- R^* و المقاسات المجوفة من نمط- R^*

مصطفى محمد جاسم*, وسن خالد

قسم الرياضيات, كلية العلوم, جامعة بغداد, بغداد, العراق

الخلاصة

الهدف من هذا البحث هو اعطاء مفاهيم جديدة للمقاس M على الحلقة , المقاسات الجزئية من هذه المفاهيم هي المقاس الجزئي الصغير من النمط- R^* وكذلك المقاس المجوف من النمط- R^* والتي تعتبر تعميمات لمفاهيم سابقه هي المقاس الجزئي الصغير والمقاس المجوف على التوالي. برهنا خواص اساسيه لهذه المفاهيم منها : صورته المقاس الجزئي الصغير من النمط- R^* تكون مقاس جزئي من النمط- R^*

1. Introduction

Let M be a unitary left R -module, and let R be any ring with one. A submodule A of M is referred to as small in M (indicated by $A \ll M$) if whenever $A + B = M$ for some $B \subseteq M$ implies that $B = M$ [1-4]. An R -module with a value greater than zero M is said to be hollow if each suitable submodule of M is small in M [5-9]. For every R -endomorphism f of M a submodule A of M if $f(A) \subseteq A$, then f is called fully invariant [10]. The cosingular submodule of M was released by Oscan in [11] in the following manner: $Z^*(M) = \{m \in M; Rm \ll E(M)\}$, where $E(M)$ is an injective hull of M see [12] and [13], where $Z^*(M)$ was called the cosingular submodule. Now, if $Z^*(M) = M$, then M is called cosingular. Many authors present generalizations of small submodule see [14-16]. Baanon and Khalid in [17] was introduced the e^* -essential submodule by using the concept of cosingular submodule. If $A \cap B \neq 0$, for each non-zero cosingular submodule B of M subsequently a submodule A is deemed to be e^* -essential. Also, in [17] it was introduced $Rad_e^*(M)$ as being the point where each maximal e^* -essential submodule

* Email: mustafa.md27182@gmail.com

intersects. If M does not include an e^* -essential maximal submodule, then $Rad_{e^*}(M) = M$. In fact it was proved that $Rad_{e^*}(M)$ is the sum of all e^* -small submodule of M , as well as $Rad_{e^*}(M)$ is a fully invariant submodule [17]. We use these concepts to introduce R^* -small submodule and investigate some properties. We also introduce and discuss the R^* -hollow module which is a generalization of the hollow module. We prove main properties of these concepts.

2. R^* -small submodule

In this section R^* -small submodule was presented with some properties.

Definition 2.1:

A submodule A of M is called Rad_R^* -small in M for short R^* -small submodule if whenever $M = A + B$, and $Rad_R^* \left(\frac{M}{B} \right) = \frac{M}{B}$ (briefly $R^* \left(\frac{M}{B} \right) = \frac{M}{B}$) implies that $B = M$. It will be denoted by $A \ll_{R^*} M$.

Remarks and examples 2.2:

1. It is evident that each small submodule is R^* -small the reverse, however is untrue. For example Z_6 as Z -module, since $Z_6 = \langle \bar{2} \rangle \oplus \langle \bar{3} \rangle$, and $R^* \left(\frac{Z_6}{\langle \bar{3} \rangle} \right) \cong R^*(\langle \bar{2} \rangle) = 0$ i.e., $R^* \left(\frac{Z_6}{\langle \bar{3} \rangle} \right) = \langle \bar{3} \rangle \neq \langle \bar{2} \rangle \cong \frac{Z_6}{\langle \bar{3} \rangle}$, but $\langle \bar{2} \rangle \neq \langle \bar{3} \rangle$, hence $\langle \bar{2} \rangle$ and $\langle \bar{3} \rangle$ are R^* -small submodules, but not small submodules.
2. As the Z -module Z_4 . In Z_4 , 0 and $\{0,2\}$ are R^* -small submodules according to (1).
3. Consider $M = Z \oplus Z_{p^\infty}$ as Z -module. Since $\frac{M}{Z} \cong Z_{p^\infty}$, therefore, $R^* \left(\frac{M}{Z} \right) = R^*(Z_{p^\infty}) = Z_{p^\infty}$ i.e., $R^* \left(\frac{M}{Z} \right) = \frac{M}{Z}$, but $M \neq Z$ which means that Z_{p^∞} is not R^* -small submodule in M .
4. Let $M = 2Z \oplus Z_2$ as Z -module. $R^* \left(\frac{M}{2Z} \right) = R^*(Z_2) = 0$, i.e., $R^* \left(\frac{M}{2Z} \right) = 2Z \neq Z_2 \cong \frac{M}{2Z}$. Thus Z_2 is R^* -small submodule in M .
5. Consider Z as Z -module. $Z = 2Z + 3Z$ as Z -module. Since $\frac{Z}{2Z} \cong Z_2$, hence $R^* \left(\frac{Z}{2Z} \right) = R^*(Z_2) = 0$ i.e., $R^* \left(\frac{M}{2Z} \right) = 2Z \neq Z_2 \cong \frac{M}{2Z}$. Thus $3Z$ is R^* -small submodule in Z .

We need to prove the following:

Proposition 2.3:

Assume that M has two submodules of M if $R^* \left(\frac{M}{A} \right) = \frac{M}{A}$ and $A \subseteq B \subseteq M$, then $R^* \left(\frac{M}{B} \right) = \frac{M}{B}$.

Proof.

Define a function $f : \frac{M}{A} \rightarrow \frac{M}{B}$ as follow $f(m + A) = m + B \forall m \in M$, it is clear that f is an epimorphism (proposition 6 in [3]) $f \left(R^* \left(\frac{M}{A} \right) \right)$ contained in $R^* \left(\frac{M}{B} \right)$. Hence $f \left(\frac{M}{A} \right) = \frac{M}{B} \subseteq R^* \left(\frac{M}{B} \right)$. Therefore, $R^* \left(\frac{M}{B} \right) = \frac{M}{B}$.

Corollary 2.4:

Consider A and B to be two submodules of M and let M be any R -module. If $R^* \left(\frac{M}{A} \right) = \frac{M}{A}$, then $R^* \left(\frac{M}{A+B} \right) = \frac{M}{A+B}$.

Proposition 2.5:

Suppose M be an R -module If $R^*(M) = M$ and $A \subseteq M$. Then $A \ll_{R^*} M$ iff $A \ll M$.

Proof.

\Rightarrow) Suppose $A + B = M, B \subseteq M$. To prove $B = M$ we claim that $R^*\left(\frac{M}{B}\right) = \frac{M}{B}$, let $f: M \rightarrow \frac{M}{B}$ be the natural epimorphism (Proposition 6 [17]) $f(R^*(M))$ contained in $R^*\left(\frac{M}{B}\right)$ and $R^*(M) = M$, then $f(M) \subseteq R^*\left(\frac{M}{B}\right)$, therefore $\frac{M}{B} \subseteq R^*\left(\frac{M}{B}\right)$ and $R^*\left(\frac{M}{B}\right) = \frac{M}{B}$, since $A \ll_{R^*} M$, then $B = M$, hence $A \ll M$.

\Leftarrow) Clearly by Remarks and examples 2.2.

These are some characteristics of R^* -small submodules.

Proposition 2.6:

1. If $A \leq B \leq M$, then $B \ll_{R^*} M$ if and only if $A \ll_{R^*} M$ and $\frac{B}{A} \ll_{R^*} \frac{M}{A}$.
2. If A and B are M submodules, then $A + B \ll_{R^*} M$ if and only if both $A \ll_{R^*} M$ and $B \ll_{R^*} M$.
3. Let $A \leq B \leq M$, if $A \leq_{R^*} B$, then $A \leq_{R^*} M$.
4. Let $f: M \rightarrow M'$ be R -homomorphism such that $A \ll_{R^*} M$, then $f(A) \ll_{R^*} M'$.
5. Let $M = M_1 \oplus M_2, A_1 \leq M_1$, and $A_2 \leq M_2$, then $A_1 \oplus A_2 \ll_{R^*} M_1 \oplus M_2$ if and only if $A_1 \ll_{R^*} M_1$ and $A_2 \ll_{R^*} M_2$.

Proof.

(1) \Rightarrow) Assume that L contained in M , in such a way that $L + A = M$ and $R^*\left(\frac{M}{L}\right) = \frac{M}{L}$, then $L + B = M$. Since $B \ll_{R^*} M$, then $L = M$. Now, let $\frac{K}{A}$ be any sub module of $\frac{M}{A}$ such that $\frac{K}{A} + \frac{B}{A} = \frac{M}{A}$ and $R^*\left(\frac{M}{K}\right) = \frac{M}{K}$, then $B + K = M$ since $B \ll_{R^*} M$, so $K = M$. Thus $\frac{K}{A} = \frac{M}{A}$.

\Leftarrow) $B + K = M, K$ contained in M , such that $R^*\left(\frac{M}{K}\right) = \frac{M}{K}$. Since $\frac{B+K}{A} = \frac{M}{A}$, so $\frac{B}{A} + \frac{A+K}{A} = \frac{M}{A}$ and $R^*\left(\frac{\frac{M}{A}}{\frac{K+A}{A}}\right) = R^*\left(\frac{M}{K+A}\right) = \frac{M}{K+A}$ by using Corollary 2.4, but $\frac{B}{A} \ll_{R^*} \frac{M}{A}$ which implies that $\frac{M}{A} = \frac{K+A}{A}$, hence $M = A + K$ as $A \ll_{R^*} M$, therefore, $K = M$.

2) Let C be a module under M such that $A + C = M$ and $R^*\left(\frac{M}{C}\right) = \frac{M}{C}$, then $A + B + C = M$ but $R^*\left(\frac{M}{C}\right) = \frac{M}{C}$ and since $A + B \ll_{R^*} M$ which implies that $C = M$, hence $A \ll_{R^*} M$. Also, $B \ll_{R^*} M$ by the same argument.

Conversely, let $A + B + C = M$ and $R^*\left(\frac{M}{C}\right) = \frac{M}{C}$. So, $A + (B + C) = M$ and $R^*\left(\frac{M}{B+C}\right) = \frac{M}{B+C}$ by using Corollary 2.4 and since $A \ll_{R^*} M$, therefore, $M = B + C$, as $B \ll_{R^*} M$ and $R^*\left(\frac{M}{C}\right) = \frac{M}{C}$, then $M = C$.

(3) Suppose that $A + K = M$ and $R^*\left(\frac{M}{K}\right) = \frac{M}{K}$, $B = M \cap B = (A + K) \cap B = (K \cap B) + A = B$ (by Modular Law), to show $R^*\left(\frac{B}{K \cap B}\right) = \frac{B}{K \cap B}$ (by Second Isomorphic Theorem) $\frac{B}{K \cap B} \cong \frac{K+B}{K} \cong \frac{M}{K}$, but $R^*\left(\frac{M}{K}\right) = \frac{M}{K}$, hence $R^*\left(\frac{B}{K \cap B}\right) = \frac{B}{K \cap B}$ and since $A \ll_{R^*} B$, then $K \cap B = B$, so $B \subseteq K$, implies that $A \subseteq K$, then $K = M$.

(4) Suppose that $f(A) + B = f(M)$, for $B \subseteq f(M)$ and $R^*\left(\frac{f(M)}{B}\right) = \frac{f(M)}{B}$. Then $A + f^{-1}(B) = M$, to show $R^*\left(\frac{M}{f^{-1}(B)}\right) = \frac{M}{f^{-1}(B)}$. Define the map $g: \frac{M}{f^{-1}(B)} \rightarrow \frac{f(M)}{B}$ by $g(x + f^{-1}(B)) = f(x) + B, \forall x \in M$ g is well defined, let $x_1 + f^{-1}(B) = x_2 + f^{-1}(B)$ iff $x_1 - x_2 \in f^{-1}(B)$ if and only if $f(x_1) - f(x_2) \in B$ if and only if $f(x_1) + B = f(x_2) + B$, g is onto, let $w \in \frac{f(M)}{B}$ so $w = f(x) + B, x \in M$, then $g(x + f^{-1}(B)) = f(x) + B$, therefore, g is onto. g is one to one, let $f(x_1) + B = f(x_2) + B$ if and only if $f(x_1) - f(x_2) \in B$ if and only if $f(x_1 - x_2) \in B$ if and only if $x_1 - x_2 \in f^{-1}(B)$ implies that $x_1 + f^{-1}(B) = x_2 + f^{-1}(B)$, hence g is isomorphism. Now, by [17] $g(R^*\left(\frac{M}{f^{-1}(B)}\right)) = R^*\left(\frac{f(M)}{B}\right) = \frac{f(M)}{B}$, therefore, $R^*\left(\frac{M}{f^{-1}(B)}\right) = g^{-1}\left(\frac{f(M)}{B}\right) = \frac{M}{f^{-1}(B)}$ implies $R^*\left(\frac{M}{f^{-1}(B)}\right) = \frac{M}{f^{-1}(B)}$. Since $A \ll_{R^*} M$, therefore, $M = f^{-1}(B)$ implies that $f(M) = B$.

(5) \Rightarrow Let $P_1: M_1 \oplus M_2 \rightarrow M_1$ be a projection map on M_1 , since $A_1 \oplus A_2 \ll_{R^*} M_1 \oplus M_2$ by (4), $P_1(A_1 \oplus A_2) \ll_{R^*} M_1$, therefore, $A_1 \ll_{R^*} M_1$. Similarly, $A_2 \ll_{R^*} M_2$.

\Leftarrow Let $J_1: M_1 \rightarrow M_1 \oplus M_2$ be the injection map from M_1 . Since $A \ll_{R^*} M_1$, therefore by (4) $J_1(A_1) = A_1 \oplus 0 \ll_{R^*} M_1 \oplus M_2$ and since $J_2: M_2 \rightarrow M_1 \oplus M_2$ by (2), then $A_1 \oplus 0 + 0 \oplus A_2 = A_1 \oplus A_2 \ll_{R^*} M_1 \oplus M_2$.

Proposition 2.7:

Let M be an R -module and $A \subseteq B \subseteq M$. If B is direct summand in M and $A \ll_{R^*} M$, then $A \ll_{R^*} B$.

Proof.

Let $A + L = B$, and B be a direct summand of M , then $M = B \oplus B_1$ for $B_1 \subseteq M$, so $M = A + L + B_1$, since $\frac{M}{L+B_1} = \frac{B+L+B_1}{L+B_1} \cong \frac{B}{B \cap (L+B_1)} = \frac{B}{L+(B \cap B_1)} = \frac{B}{L}$ (by using Second Isomorphism Theorem and Modular Law). Consequently, $R^*\left(\frac{B}{L}\right) = \frac{B}{L}$, then by Corollary 2.4, $R^*\left(\frac{M}{L+B_1}\right) = \frac{M}{L+B_1}$, but $A \ll_{R^*} M$, then $M = L + B_1$. Now, $B = B \cap M = B \cap (L + B_1) = L + (B \cap B_1) = L$ (by Modular Law). Thus, $A \ll_{R^*} B$.

Proposition 2.8:

Assume that M is an R -module and A, B , and C are submodules of M with $A \subseteq B \subseteq C \subseteq M$, if $B \ll_{R^*} C$, then $A \ll_{R^*} M$.

Proof.

Suppose that $A + L = M$ and $R^*\left(\frac{M}{L}\right) = \frac{M}{L}$, so $B + L = M$, since $C \subseteq M$, hence $C = M \cap C = (B + L) \cap C = B + (L \cap C)$ by Modular Law. To demonstrate $R^*\left(\frac{C}{L \cap C}\right) = \frac{C}{L \cap C}$, since $\frac{C}{L \cap C} \cong \frac{L+C}{C} \cong \frac{M}{C}$ by Second Isomorphic Theorem, but $R^*\left(\frac{M}{C}\right) = \frac{M}{C}$, then $R^*\left(\frac{C}{L \cap C}\right) = \frac{C}{L \cap C}$. Since $B \ll_{R^*} C$, then $C = L \cap C$, so $C \subseteq L$, but $A \subseteq C$. Hence, $A \subseteq L$, since $A + L = M$, then $L = M$. Therefore, $A \ll_{R^*} M$.

Remark 2.9:

It is uncommon for the converse of Proposition 2.7 to be true. The example that follows demonstrates. Consider $M = Z \oplus Z_{p^\infty}$ as Z -module $0 \oplus Z \subseteq 0 \oplus Z_{p^\infty} \subseteq Z \oplus Z_{p^\infty}$ it is clear that $Z \ll_{R^*} Z \oplus Z_{p^\infty}$, but Z_{p^∞} is not R^* -small in $Z \oplus Z_{p^\infty}$ (see 2.2 (3)).

3. R^* -hollow module

Definition 3.1:

A module M is non-zero if for each proper submodule of M is Rad_R $*$ -small submodule of M , then M referred to an Rad_R $*$ -hollow module (or simply R^* -hollow module).

Remarks and examples 3.2:

1. Every simple module is R^* -hollow module. For example, Z_p as Z -module, (where p is a prime number).
2. It is obvious that any hollow module is R^* -hollow module. However, the reverse is not true. For example, Z_6 as Z -module.
3. Z_4 as Z -module is R^* -hollow module.
4. Consider $M = Z \oplus Z_{p^\infty}$ as Z -module is not R^* -hollow module. Since Z_{p^∞} is proper submodule of M , but Z_{p^∞} is not R^* -small module of M .

The resulting theorem gives an explains of the R^* -hollow module.

Theorem 3.3:

M is R^* -hollow module if and only if for each proper submodule A of M is small in M with $R^* \left(\frac{M}{A} \right) = \frac{M}{A}$ where M is R -module.

Proof.

\Rightarrow) Assuming that A is a proper submodule of M such that $R^* \left(\frac{M}{A} \right) = \frac{M}{A}$. We have to show that $A \ll M$. Assume that there exists a proper submodule B of M such that $M = A + B$, since M is R^* -hollow, then $B \ll_{R^*} M$, but $R^* \left(\frac{M}{A} \right) = \frac{M}{A}$, then $M = A$ which is a contradiction. Thus $A \ll M$.

\Leftarrow) To demonstrate that M is R^* -hollow module. Take A is a proper submodule of M . Assume that A is not R^* -small, there exists a proper submodule B of M such that $R^* \left(\frac{M}{B} \right) = \frac{M}{B}$ and $M = A + B$. By our assumption $B \ll M$, then $A = M$ which is a contradiction. Thus, M is R^* -hollow.

Proposition 3.4:

A non-zero epimorphic image of R^* -hollow module is R^* -hollow.

Proof.

Given $f: M \rightarrow M'$ be an epimorphism and M be R^* -hollow module and assume $B \subsetneq M'$, therefore $f^{-1}(B) \subsetneq M$. To show that $f^{-1}(B)$ is a proper in M . If $f^{-1}(B) = M$, then $B = f(M) = M'$ which implies $B = M'$ and that is a contradiction, so $f^{-1}(B)$ is a proper submodule of M . Since M is an R^* -hollow, therefore $f^{-1}(B) \ll_{R^*} M$ and, by Proposition 2.6 we have $f(f^{-1}(B)) \ll_{R^*} M'$, then $B \ll_{R^*} M'$.

Corollary 3.5:

Suppose M is an R -module and $A \subseteq M$. if M is an R^* -hollow, then $\frac{M}{A}$ is an R^* -hollow.

Proof.

Let $f: M \rightarrow \frac{M}{A}$ be a natural epimorphism and M be an R^* -hollow Proposition 3.4, $\frac{M}{A}$ is an R^* -hollow.

Remark 3.6:

It is not required that the direct sum of an R^* -hollow modules be R^* -hollow. For instance $M = Z \oplus Z_{p^\infty}$ as Z -module as previously demonstrated that every proper submodule of Z is R^* -small so Z as Z -module is R^* -hollow and every proper submodule of Z_{p^∞} is not big, hence is R^* -small so Z_{p^∞} as Z -module is R^* -hollow. However, M is not an R^* -hollow.

The direct sum of R^* -hollow modules is an R^* -hollow under the criteria we will now list. Not that a submodule A of M is called completely invariant for each $f \in \text{End}(M)$, $f(A) \subseteq A$. In [11], [18], likewise a module M is said to be a duo module if for each of its submodules is completely invariant.

Proposition 3.7:

Let $M = M_1 \oplus M_2$, M be a duo module, then M is an R^* -hollow if and only if M_1 and M_2 are R^* -hollow. Provided that $A \cap M_1 \neq M_i$, for all $i = 1, 2$, $A \subseteq M$.

Proof.

\Rightarrow) Let M be an R^* -hollow module and $A_1 \oplus A_2 \subsetneq M_1 \oplus M_2$ with $A_1 \subsetneq M_1$ and $A_2 \subsetneq M_2$. Consider that $\pi_1: M_1 \oplus M_2 \rightarrow M_1$ be a projection map, which defined as follows, $\pi_1(m_1 + m_2) = m_1$, for all $m_1 \in M_1, m_2 \in M_2$, since $A_1 \oplus A_2 \ll_{R^*} M_1 \oplus M_2$. Then by Proposition 2.6, we have $\pi_1(A_1 \oplus A_2) \ll_{R^*} \pi_1(M_1 \oplus M_2)$, so $A \ll_{R^*} M_1$. That means M_1 is an R^* -hollow. By using the same way one can show that M_2 is an R^* -hollow.

\Leftarrow) Let M_1 and M_2 be an R^* -hollow, and let $A \subsetneq M$. Since M is a duo module, then $A = A_1 \oplus A_2$, where $A_1 = A \cap M_1$ and $A_2 = A \cap M_2$, since $A_1 \ll_{R^*} M_1$ and $A_2 \ll_{R^*} M_2$. Then by Proposition 2.6, $A_1 \oplus A_2 \ll_{R^*} M = M_1 \oplus M_2$.

A module M is said to be distributive if for each of its submodules A, B , and C in M such that $(A + B) \cap C = A \cap C + B \cap C$ [19], [20].

Proposition 3.8:

Assuming that $M = M_1 \oplus M_2$ be a distributive module, then M is an R^* -hollow if and only if M_1 and M_2 are R^* -hollow are provided that $A \cap M_i \neq M_i$ for all $i = 1, 2$.

Proof.

By using the same argument in Proposition 3.7.

References

- [1] F.Kash, Module and rings, Londone : Acadimic Oress, 1982.
- [2] S. Mohammed and j.Bruno.Muller, Continous and discret modules, New york:Lens147: Cambridge Press, 1990.
- [3] W. Frank and R.Kent, Rings and categories of module, verlag: springer, 1994.
- [4] Y. Zhou, "Generalizations of perfect, Semiperfect,and semiregular rings," *Algebra Colloquium*, vol. 7, no. 3, pp. 305-318, 2000.
- [5] P.Fleury, "Hollow modules and local endomorphism rings," *Pacific journal of mathematics*, vol. 2, no. 53, pp. 379-385, 1974.
- [6] I.Mohammed, Module theory, Baghdad: bookstore for printing publishing and translating, 2021.
- [7] j. Clark, C. Iomp, N. Vanaja and R. Wisbauer, Lifting Module Supplement and Projectivity in Module Theory, Berlline, 2006.
- [8] A.Azizi, "Hollow module over commutative Ring," *Palestine journal of mathmatica*, vol. 3, no. spec1, pp. 449-456, 2014.

- [9] N. Orhan, T. D. Keskin and R. tribak, "On hollow- lifting module," *Taiwanese journal of mathematics*, vol. 11, no. 2, pp. 545-568, 2007.
- [10] C.Ozcan, "Duo modules," *Glasgow mathematical journal trust*, vol. 48, pp. 533-545, 2006.
- [11] A.Ozcan, "Modules with small cyclic sub modules in their injective hulls," *Taylor and francis online* , vol. 30, pp. 1575-1589, 2002.
- [12] K.R.Goodearl, *Ring theory NonSingular Rings and Modules*, Salt Lake,Utah: Marcel Dekker,Inc, 1976.
- [13] K.R.Goodearl and R.B.warfield, *An Intoducton to Non commutative Noetherain rings*, Cambridge: Cambridge university press, 2004.
- [14] M.Abbas and F. Mohamed, "On e-smallsubmodules," *Ibn-Alhatham journal for pure and apply.sci*, vol. 28, pp. 214-222, 2015.
- [15] A.Kabban and W.Kalid, "On jacobson-small submodules," *Iraqi journal of science*, vol. 30, pp. 1584-1591, 2019.
- [16] W.Khalid and M.K.Enaas, "On generalization of smallsubmodules," *sci.int.(lahore)*, vol. 30, pp. 359-366, 2018.
- [17] H.Baanoon and W.Khalid, "e*-essential small submodules and e*-hollow modules," *European journal of apure and applide mathematics*, vol. 15, pp. 478-485, 2022.
- [18] I.Mohammed and N. Rana, "On P-Duo Module," *International Journal of Algebra*, vol. 8, no. 5, pp. 229-238, 2014.
- [19] V.Camillo, "Distributive modules," *Journal of algebra* , vol. 36, pp. 16-25, 1975.
- [20] V.Erdogdu, "Distributive modules," *Canada.Math.Bull*, vol. 2, pp. 248-254, 1987.