



ISSN: 0067-2904

Cubic Trigonometric Spline for Solving Nonlinear Volterra Integral Equations

Nabaa N.Hasan, Majid R.Nasif*

Department of Mathematics, College of Science, University of Mustansiriyah, Baghdad, Iraq

Received: 4/5/ 2019

Accepted: 17/ 7/2019

Abstract

In this paper, cubic trigonometric spline is used to solve nonlinear Volterra integral equations of second kind. Examples are illustrated to show the presented method's efficiency and convenience.

Keywords: Nonlinear integral equation, Volterra second kind , Cubic trigonometric spline.

شريحة مثلثية مكعبية لحل معادلات فولتيرا التكاملية غير الخطية

نبا حسن* ، ماجد ناصيف

قسم الرياضيات ، كلية العلوم ، الجامعة المستنصرية ، بغداد ، العراق

الخلاصة

في هذا البحث ، يتم استخدام شريحة مثلثية مكعب لحل معادلات فولتيرا التكاملية غير الخطية من النوع الثاني. تم توضيح أمثلة لإظهار كفاءة الطريقة المعروضة وراحتها.

1. Introduction

An integral equation is defined as an equation in which the unknown function $u(x)$ to be determined appear under the integral sign. The subject of integral equations is one of the most useful mathematical tools in both pure and applied mathematics. It has enormous applications in many physical problems. Many initial and boundary value problems associated with ordinary differential equation (ODE) and partial differential equation (PDE) can be transformed into problems of solving some approximate integral equations.

The importance of trigonometric functions in different areas, such as electronics or medicine is well known. Recently, trigonometric splines and polynomials gained very high interest with computer-aided geometric design. In particular, curve designs [1, 2, 3] construct a cubic trigonometric Bezier curve with two shape parameters, on the basis of cubic trigonometric Bernstein functions to generate a curve interpolation scheme.

The solution of integral equation can be approximated by using the non-polynomial spline functions and the collection method [4]. Combination of affixed point method and cubic B-spline functions was used [5] to solve the integral equation numerically. The error analysis for the method shows that the approximate solution converges to the exact solution. Non-polynomial spline of functions was used [6] to develop numerical methods to approximate the solution of second kind Volterra integral equations.

There is another method to solve these problems, which is called the Laplace transform series decomposition method (LTSDM), but the cubic trigonometric spline method is the best method

*Email: alzaer1972@uomustansiriyah.edu.iq

because it gives approximate solution which is very close to the exact solution, as shown in the tables that will be presented in this paper.

In this paper, the presentation of the algorithm is achieved in a very simplified way so that the reader can apply it to other types of equations such as Fredholm integral equations.

Previous studies provided more details on trigonometric spline [7, 8, 9, 10].

2. Trigonometric cubic spline method:-[7]

In a simple way, we take [a,b] as interval, in order to improve the numerical method for approximation solution of the following kind:

$$u(x) = f(x) + \int_0^x k(x,t,u(t)) dt \quad \dots (1)$$

For this purpose define $x_j = a + j h, j = 0,1,2, \dots, L$

For all j th segment, the cubic non-polynomial spline $S_j(x)$ has the form:-

$$S_j(x) = a_j \sin k(x - x_j) + b_j \cos k(x - x_j) + c_j (x - x_j) + d_j \quad \dots (2)$$

Where a_j, b_j, c_j and d_j are real finite constants and k is the frequency of the trigonometric function which will be used to develop the accuracy of this method. In this paper we consider $k \in (0, 1]$, where the value of k can be determined by mathematical programs such as Matlab14. or Mathcad15, by repeating values for k until we reach the random value that gives the best approximate solution.

Now, we will explain this approximate method by considering the following relation:

$$S'_j(x_j) = a_j k + c_j \approx u'(x_j)$$

$$S''_j(x_j) = -b_j k^2 \approx u''(x_j)$$

$$S'''_j(x_j) = -a_j k^3 \approx u'''(x_j)$$

Now, we can obtain the values of a_j, b_j, c_j and d_j as follows:

$$a_j = -\frac{1}{k^3} u'''(x_j) \quad \dots (3)$$

$$b_j = -\frac{1}{k^2} u''(x_j) \quad \dots (4)$$

$$c_j = u'(x_j) + \frac{1}{k^2} u'''(x_j) \quad \dots (5)$$

$$d_j = u(x_j) + \frac{1}{k} u''(x_j) \quad \dots (6)$$

Now, we differentiate equation (1) three times with respect to x and then put:

$x = x_j, j = 0,1,2, \dots, L - 1$ and for clarification we will compensate $x_j = 0$

and in the same way for other values:

$$u(0) = f(0) \quad \dots (7)$$

$$u'(x) = f'(x) + \int_0^x \frac{\partial k(x,t,u(t))}{\partial x} dt + k(x,x,u(x))$$

$$u'(0) = f'(0) + k(0,0,u(0)) \quad \dots (8)$$

$$\text{Let } E(x,t,u(t)) = \frac{\partial k(x,t,u(t))}{\partial x}$$

$$u''(x) = f''(x) + \int_0^x \frac{\partial E(x,t,u(t))}{\partial x} dt + 2E(x,x,u(x))$$

$$u''(0) = f''(0) + 2E(0,0,u(0)) \quad \dots (9)$$

$$\text{Let } F(x,t,u(t)) = \frac{\partial E(x,t,u(t))}{\partial x}$$

$$u'''(x) = f'''(x) + \int_0^x \frac{\partial F(x,t,u(t))}{\partial x} dt + 3F(x,x,u(x))$$

$$u'''(0) = f'''(0) + 3F(0,0,u(0)) \quad \dots (10)$$

Thus, we can approximate the solution of nonlinear Volterra integral equations of second kind (1) by using equation (2).

Algorithm:

Step1:- Input $h = \frac{b-a}{L}, x_j = x_0 + j h, j = 0,1,2, \dots, L, x_0 = a, x_L = b$

Step2:- Compute a_j, b_j, c_j and d_j by substituting the equations 7 – 10 in equations 3 – 6

Step3:- Evaluate $S_0(x)$ by using Step2 and equation (2) for $j = 0$

Step4:- Approximate $u_1 = u(x_1) \approx S_0(x_1)$

Step5:- Do the following steps for $j = 1$ to $L - 1$

Step6:- Compute a_j, b_j, c_j and d_j by using equation (3 – 6) and replacing $u'(x_j), u''(x_j)$ and $u'''(x_j)$ in $S_j^i(x_j), S_j^{ii}(x_j)$ and $S_j^{iii}(x_j)$

Step7:- Approximate $u_{j+1} = S_j(x_{j+1})$

To show that the presented method is efficient and convenient:

Example (1):- Consider the following non-linear Volterra integral equation of second kind:-

$$u(x) = \sin(x) + \frac{2}{3} \cos(x) - \frac{1}{3} \cos^2(x) - \frac{1}{3} + \int_0^x \sin(x-t) u^2(t) dt$$

Which has the exact solution $u(x) = \sin(x)$, [8]

If $h = 0.1$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0.0 \\ S_{10}(x) & \text{if } 0.0 \leq x \leq 0.1 \\ S_{11}(x) & \text{if } 0.1 \leq x \leq 0.2 \\ S_{12}(x) & \text{if } 0.2 \leq x \leq 0.3 \\ S_{13}(x) & \text{if } 0.3 \leq x \leq 0.4 \\ S_{14}(x) & \text{if } 0.4 \leq x \leq 0.5 \\ S_{15}(x) & \text{if } 0.5 \leq x \leq 0.6 \\ S_{16}(x) & \text{if } 0.6 \leq x \leq 0.7 \\ S_{17}(x) & \text{if } 0.7 \leq x \leq 0.8 \\ S_{18}(x) & \text{if } 0.8 \leq x \leq 0.9 \\ S_{19}(x) & \text{if } 0.9 \leq x \leq 1.0 \end{cases}$$

Table 1- Approximate Results for Example 1 if $h = 0.1$

X_j	Exact solution	Approximate solution	Absolute error
0.0	0.0	0.0	0.0
0.1	0.099833416646828	0.099833416646828	0.0
0.2	0.19866933079506	0.19866933079506	0.0
0.3	0.29552020666134	0.29552020666134	0.0
0.4	0.38941834230865	0.38941834230865	0.0
0.5	0.4794255386042	0.4794255386042	0.0
0.6	0.56464247339504	0.56464247339503	0.000000000000001
0.7	0.64421768723769	0.64421768723768	0.000000000000001
0.8	0.71735609089952	0.71735609089952	0.0
0.9	0.78332690962748	0.78332690962748	0.0
1.0	0.8414709848079	0.84147098480789	0.000000000000001
L.S.E.		2.2325351775545e-28	




Figure 1-Comparison of Exact Solution and Approximate Solution if $h = 0.1$

If $h = \frac{1}{3}$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0 \\ S_{10}(x) & \text{if } 0 \leq x \leq \frac{1}{3} \\ S_{11}(x) & \text{if } \frac{1}{3} \leq x \leq \frac{2}{3} \\ S_{12}(x) & \text{if } \frac{2}{3} \leq x \leq 1 \end{cases}$$

Table 2-Approximate Results for Example 1 if $h = (1)/3$

x_j	Exact solution	Approximate solution	Absolute error
0.0	0.0	0.0	0.0
$\frac{1}{3}$	0.32719469679615	0.32719469679615	0.0
$\frac{2}{3}$	0.61836980306974	0.61836980306974	0.0
1.0	0.8414709848079	0.8414709848079	0.0
L.S.E.		9.6481363554251e-29	




Figure 2-Comparison of Exact Solution and Approximate Solution if $h = (1/3)$

Example(2):-Consider the following non-linear Volterra integral equation of second kind:-

$$u(x) = x - x e^x + \int_0^x e^{x-t} e^{u(t)} dt$$

Which has the exact solution $u(x) = x$,[8]

If $h = 0.1$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0.0 \\ S_{10}(x) & \text{if } 0.0 \leq x \leq 0.1 \\ S_{11}(x) & \text{if } 0.1 \leq x \leq 0.2 \\ S_{12}(x) & \text{if } 0.2 \leq x \leq 0.3 \\ S_{13}(x) & \text{if } 0.3 \leq x \leq 0.4 \\ S_{14}(x) & \text{if } 0.4 \leq x \leq 0.5 \\ S_{15}(x) & \text{if } 0.5 \leq x \leq 0.6 \\ S_{16}(x) & \text{if } 0.6 \leq x \leq 0.7 \\ S_{17}(x) & \text{if } 0.7 \leq x \leq 0.8 \\ S_{18}(x) & \text{if } 0.8 \leq x \leq 0.9 \\ S_{19}(x) & \text{if } 0.9 \leq x \leq 1.0 \end{cases}$$

Table 3-Approximate Results for Example 2 if $h = 0.1$

X_i	Exact solution	Approximate solution	Absolute error
0.0	0.0	0.0	0.0
0.1	0.1	0.1	0.0
0.2	0.2	0.2	0.0
0.3	0.3	0.3	0.0
0.4	0.4	0.4	0.0
0.5	0.5	0.5	0.0
0.6	0.6	0.6	0.0
0.7	0.7	0.7	0.0
0.8	0.8	0.8	0.0
0.9	0.9	0.9	0.0
1.0	1.0	1.0	0.0
L.S.E.		0.0	




Figure 3-Comparison of Exact Solution and Approximate Solution if $h = 0.1$

If $h = \frac{1}{3}$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0 \\ S_{10}(x) & \text{if } 0 \leq x \leq \frac{1}{3} \\ S_{11}(x) & \text{if } \frac{1}{3} \leq x \leq \frac{2}{3} \\ S_{12}(x) & \text{if } \frac{2}{3} \leq x \leq 1 \end{cases}$$

Table 4-Approximate Results for Example 2 if $h = (1)/3$

X_j	Exact solution	Approximate solution	Absolute error
0.0	0.0	0.0	0.0
$\frac{1}{3}$	$\frac{1}{3}$	0.333333333333333	3.33333333333333e-15
$\frac{2}{3}$	$\frac{2}{3}$	0.666666666666666	6.66666666666667e-15
1	1	0.999999999999999	1.0e-14
L.S.E.		1.55555555555556e-28	




Figure 4-Comparison of Exact Solution and Approximate Solution if $h = (1)/3$

Example(3):-Consider the following non-linear Volterra integral equation of second kind: $-u(x) = \cos(x) - \frac{1}{2}x^2 + \int_0^x (x-t)(u^2(t) + \sin^2(t)) dt$

Which has the exact solution $u(x) = \cos(x)$, [7]

$$h = \frac{1}{3}$$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0 \\ S_{10}(x) & \text{if } 0 \leq x \leq \frac{1}{3} \\ S_{11}(x) & \text{if } \frac{1}{3} \leq x \leq \frac{2}{3} \\ S_{12}(x) & \text{if } \frac{2}{3} \leq x \leq 1 \end{cases}$$

Table 5-Approximate Results for Example 3 if $h = \frac{1}{3}$

X_j	Exact solution	Approximate solution	Absolute error
0.0	1.0	1.0	0.0
$\frac{1}{3}$	0.94495694631474	0.94495694631474	0.0
$\frac{2}{3}$	0.78588726077695	0.78588726077695	0.0
1	0.54030230586814	0.54030230586814	0.0
L.S.E.		9.5314704530475e-30	




Figure 5 -Comparison of Exact Solution and Approximate Solution if $h = (1/3)$

$$h = 0.1$$

$$S(x) = \begin{cases} 0 & \text{if } -1 \leq x \leq 0.0 \\ S_{10}(x) & \text{if } 0.0 \leq x \leq 0.1 \\ S_{11}(x) & \text{if } 0.1 \leq x \leq 0.2 \\ S_{12}(x) & \text{if } 0.2 \leq x \leq 0.3 \\ S_{13}(x) & \text{if } 0.3 \leq x \leq 0.4 \\ S_{14}(x) & \text{if } 0.4 \leq x \leq 0.5 \\ S_{15}(x) & \text{if } 0.5 \leq x \leq 0.6 \\ S_{16}(x) & \text{if } 0.6 \leq x \leq 0.7 \\ S_{17}(x) & \text{if } 0.7 \leq x \leq 0.8 \\ S_{18}(x) & \text{if } 0.8 \leq x \leq 0.9 \\ S_{19}(x) & \text{if } 0.9 \leq x \leq 1.0 \end{cases}$$

Table 6-Approximate Results for Example 3 if $h = 0.1$

X_j	Exact solution	Approximate solution	Absolute error
0.0	1.0	1.0	0.0
0.1	0.99500416527803	0.99500416527803	0.0
0.2	0.98006657784124	0.98006657784125	0.00000000000001
0.3	0.95533648912561	0.95533648912562	0.00000000000001
0.4	0.92106099400289	0.92106099400289	0.0
0.5	0.87758256189037	0.87758256189038	0.00000000000001
0.6	0.82533561490968	0.82533561490969	0.00000000000001
0.7	0.76484218728449	0.7648421872845	0.00000000000001
0.8	0.69670670934717	0.69670670934718	0.00000000000001
0.9	0.62160996827066	0.62160996827068	0.00000000000002
1.0	0.54030230586814	0.54030230586816	0.00000000000002
L.S.E.		1.4970818729916e-27	




Figure 6 -Comparison of Exact Solution and Approximate Solution if $h=0.1$

3. Conclusion

The present work is an effort to obtain the approximate solution of Volterra integral equation of the second kind by using trigonometric cubic spline method. Three test examples were considered with the exact solution, for which the results are given in Tables-(1, 2, 3, 4, and 5) and Figures-(1, 2, 3, 4, and 5) to show the accuracy and efficiency of the method. All the computations were performed by Mathcad 15

References

1. Huayong, L. and Daming, Z. **2012**. "Cubic Trigonometric Polynomial B-spline Curves and Surfaces with Shape Parameter", *Journal of Information & Computation of Science*, **9**(4): 989-996.
2. Uzma, B. and Jamaludin, A. **2014**. "A Rational Trigonometric Spline to Visualize Positive Data", AIP conference proceedings 1605, 286.
3. Muhammad, S. and Nabila, J. **2017**. "A Rational Quadratic Trigonometric Spline With Interval Shape Control" 14th International conference on computer Graphics, Imaging and visualization, 23 -25 May, 2017.
4. Maleknejad, Kh. and Jalilian, H. **2018**. "Non-polynomial Spline Functions and Quasi – linearization to Approximate Nonlinear Volterra Integral Equation", Faculty of sciences and Mathematics university of Nis, Serbia, pp.3947-3956, 2018.
5. Khosrow, M. and Reza, M. **2014**. "Numerical solution of Volterra functional integral equation by using cubic B-spline scaling functions", Wiley periodicals, Inc. Numerical Methods partial Differential Eq, **30**: 699-722.
6. Muna, M. and Sarah, H. **2014**. "Solution of Second Kind Volterra Integral Equations Using Non-Polynomial Spline Functions", *Baghdad Science Journal*, **11**(2): 406-41.
7. Borhan, F. and Abbas, H. **2016**. "Non-Polynomial Spline Method for the Solution of the System of two Nonlinear Volterra Integral Equations", *Kirkuk university Journal*, **11**(3): 15-25.
8. Wazwaz, A.M. **2011**. "Linear and Nonlinear Integral Equation", Methods and Applications, Springer.
9. Hossin, P.A. **2012**. "The Solve of Integral Differential Equation by Non-Polynomial Spline Function and Quadrature Formula", International Conference on Applied mathematics and Pharmaceutical Science, Jan. 7-8, P.595-597, 2012.
10. Saba, N.M. **2014**. "Solution of Second Kind Volterra Integral Equations using Linear Non-polynomial Spline Function" *Mathematical Theory and Modeling*, **4**(9): 2014.