



On Nano Generalized Semi Generalized Closed Sets

Qays Hatem Imran*

Department of Mathematics, College of Education for Pure Science, Al-Muthanna University, Samawah, Iraq

Abstract:

In this paper we introduced a new class of N-CS called $Ngsg$ -CS and study their basic properties in nano topological spaces. We also introduce $Ngsg$ -closure and $Ngsg$ -interior and study some of their fundamental properties.

Keywords: $Ngsg$ -CS, $Ngsg$ -OS, $Ngsg$ -closure and $Ngsg$ -interior.

مجموعات النانو المعممة شبه المعممة المغلقة

قيس حاتم عمران*

قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة المثنى، السماوة، العراق

الخلاصة:

في هذا البحث قدمنا فئة جديدة من مجموعات النانو المغلقة تسمى بمجموعات النانو المعممة شبه المعممة المغلقة و دراسة خصائصها الأساسية في الفضاءات النانو التولوجية. قدمنا أيضا انغلاق النانو المعممة شبه المعممة و مجموعة النقاط الداخلية النانو المعممة شبه المعممة و دراسة بعض خصائصها الأساسية.

1. Introduction

M. Lellis Thivagar and Carmel Richard [1] introduced nano topological space (or simply NTS) with respect to a subset X of a universe which is defined in terms of lower and upper approximations of X . He has also defined nano closed sets (briefly N-CS), nano interior and nano closure of a set. In 2014, Ng -CS was introduced by K. Bhuvanewari and K. Mythili Gnanapriya [2]. K. Bhuvanewari and A. Ezhilarasi [3] introduced the concept of Nsg -CS and Ng -CS in NTS. The concept gsg -CS have been introduced and studied by M. Lellis et al [4] in classical topology. The purpose of this paper is to introduce the concept of $Ngsg$ -CS and study their basic properties in NTS. We also introduce $Ngsg$ -closure and $Ngsg$ -interior and obtain some of its properties.

2. Preliminaries

Throughout this paper, $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and $(\mathcal{V}, \sigma_{\mathcal{R}}(Y))$ (or simply \mathcal{U} and \mathcal{V}) always mean NTS on which no separation axioms are assumed unless otherwise mentioned. For a set \mathcal{A} in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, $Ncl(\mathcal{A})$, $Nint(\mathcal{A})$ and $\mathcal{A}^c = \mathcal{U} - \mathcal{A}$ denote the nano closure of \mathcal{A} , the nano interior of \mathcal{A} and the nano complement of \mathcal{A} respectively.

Definition 2.1:[5] Let \mathcal{U} be a non-empty finite set of objects called the universe and \mathcal{R} be an equivalence relation on \mathcal{U} named as the indiscernibility relation. Elements belonging to the same equivalence class are said to be indiscernible with in another. The pair $(\mathcal{U}, \mathcal{R})$ is called the approximation space.

Remark 2.2:[5] Let $(\mathcal{U}, \mathcal{R})$ be an approximation space and $X \subseteq \mathcal{U}$. Then:

- i. The lower approximation of X with respect to \mathcal{R} is the set of all objects, which can be for certain classified as X with respect to \mathcal{R} and it is denoted by $L_{\mathcal{R}}(X)$. That is, $L_{\mathcal{R}}(X) = \cup\{\mathcal{R}(x): \mathcal{R}(x) \subseteq X, x \in \mathcal{U}\}$, where $\mathcal{R}(x)$ denotes the equivalence class determined by x .

*Email: qays.imran@mu.edu.iq

- ii. The upper approximation of X with respect to \mathcal{R} is the set of all objects, which can be possibly classified as X with respect to \mathcal{R} and it is denoted by $U_{\mathcal{R}}(X)$. That is, $U_{\mathcal{R}}(X) = \cup\{\mathcal{R}(x): \mathcal{R}(x) \cap X \neq \phi, x \in \mathcal{U}\}$.
- iii. The boundary region of X with respect to \mathcal{R} is the set of all objects, which can be classified neither as X nor as not X with respect to \mathcal{R} and it is denoted by $B_{\mathcal{R}}(X)$. That is, $B_{\mathcal{R}}(X) = U_{\mathcal{R}}(X) - L_{\mathcal{R}}(X)$.

Proposition 2.3:[6] If $(\mathcal{U}, \mathcal{R})$ is an approximation space and $X, Y \subseteq \mathcal{U}$. Then:

- i. $L_{\mathcal{R}}(X) \subseteq X \subseteq U_{\mathcal{R}}(X)$.
- ii. $L_{\mathcal{R}}(\phi) = U_{\mathcal{R}}(\phi) = \phi$ and $L_{\mathcal{R}}(\mathcal{U}) = U_{\mathcal{R}}(\mathcal{U}) = \mathcal{U}$.
- iii. $U_{\mathcal{R}}(X \cup Y) = U_{\mathcal{R}}(X) \cup U_{\mathcal{R}}(Y)$.
- iv. $U_{\mathcal{R}}(X \cap Y) \subseteq U_{\mathcal{R}}(X) \cap U_{\mathcal{R}}(Y)$.
- v. $L_{\mathcal{R}}(X \cup Y) \supseteq L_{\mathcal{R}}(X) \cup L_{\mathcal{R}}(Y)$.
- vi. $L_{\mathcal{R}}(X \cap Y) = L_{\mathcal{R}}(X) \cap L_{\mathcal{R}}(Y)$.
- vii. $L_{\mathcal{R}}(X) \subseteq L_{\mathcal{R}}(Y)$ and $U_{\mathcal{R}}(X) \subseteq U_{\mathcal{R}}(Y)$ whenever $X \subseteq Y$.
- viii. $U_{\mathcal{R}}(X^c) = (L_{\mathcal{R}}(X))^c$ and $L_{\mathcal{R}}(X^c) = (U_{\mathcal{R}}(X))^c$.
- ix. $U_{\mathcal{R}}U_{\mathcal{R}}(X) = L_{\mathcal{R}}U_{\mathcal{R}}(X) = U_{\mathcal{R}}(X)$.
- x. $L_{\mathcal{R}}L_{\mathcal{R}}(X) = U_{\mathcal{R}}L_{\mathcal{R}}(X) = L_{\mathcal{R}}(X)$.

Definition 2.4:[1] Let \mathcal{U} be the universe, \mathcal{R} be an equivalence relation on \mathcal{U} and $\tau_{\mathcal{R}}(X) = \{\phi, \mathcal{U}, L_{\mathcal{R}}(X), U_{\mathcal{R}}(X), B_{\mathcal{R}}(X)\}$ where $X \subseteq \mathcal{U}$. Then by proposition (2.3), $\tau_{\mathcal{R}}(X)$ satisfies the following axioms:

- i. $\phi, \mathcal{U} \in \tau_{\mathcal{R}}(X)$.
- ii. The union of the elements of any subcollection of $\tau_{\mathcal{R}}(X)$ is in $\tau_{\mathcal{R}}(X)$.
- iii. The intersection of the elements of any finite subcollection of $\tau_{\mathcal{R}}(X)$ is in $\tau_{\mathcal{R}}(X)$.

That is, $\tau_{\mathcal{R}}(X)$ is a topology on \mathcal{U} called the nano topology on \mathcal{U} with respect to X and the pair $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is called a nano topological space (or simply NTS). The elements of $\tau_{\mathcal{R}}(X)$ are called as nano open sets (briefly N-OS).

Remark 2.5:[1] Let $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ be a NTS with respect to X where $X \subseteq \mathcal{U}$ and \mathcal{R} be an equivalence relation on \mathcal{U} . Then \mathcal{U}/\mathcal{R} denotes the family of equivalence classes of \mathcal{U} by \mathcal{R} .

Definition 2.6:[1] A subset \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be:

- i. a nano semi-open set (briefly Ns-OS) if $\mathcal{A} \subseteq Ncl(Nint(\mathcal{A}))$ and a nano semi-closed set (briefly Ns-CS) if $Nint(Ncl(\mathcal{A})) \subseteq \mathcal{A}$. The nano semi-closure of a set \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is the intersection of all Ns-CS that contain \mathcal{A} and is denoted by $Nscl(\mathcal{A})$.
- ii. a nano α -open set (briefly $N\alpha$ -OS) if $\mathcal{A} \subseteq Nint(Ncl(Nint(\mathcal{A})))$ and a nano α -closed set (briefly $N\alpha$ -CS) if $Ncl(Nint(Ncl(\mathcal{A}))) \subseteq \mathcal{A}$. The nano α -closure of a set \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is the intersection of all $N\alpha$ -CS that contain \mathcal{A} and is denoted by $Nacl(\mathcal{A})$.

Definition 2.7: A subset \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be:

- i. a nano generalized closed set (briefly Ng-CS) [2] if $Ncl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a N-OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The complement of a Ng-CS is a Ng-OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.
- ii. a nano αg -closed set (briefly $N\alpha g$ -CS) [7] if $Nacl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a N-OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The complement of a $N\alpha g$ -CS is a $N\alpha g$ -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.
- iii. a nano $g\alpha$ -closed set (briefly $Ng\alpha$ -CS) [7] if $Nacl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a $N\alpha$ -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The complement of a $Ng\alpha$ -CS is a $Ng\alpha$ -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.
- iv. a nano sg -closed set (briefly Nsg -CS) [3] if $Nscl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a Ns-OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The complement of a Nsg -CS is a Nsg -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.
- v. a nano gs -closed set (briefly Ngs -CS) [3] if $Nscl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a N-OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The complement of a Ngs -CS is a Ngs -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proposition 2.8:[1,2] In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then the following statements hold and the converse of each statements are not true:

- i. Every N-OS (resp. N-CS) is a $N\alpha$ -OS (resp. $N\alpha$ -CS).
- ii. Every N-OS (resp. N-CS) is a Ng-OS (resp. Ng-CS).
- iii. Every $N\alpha$ -OS (resp. $N\alpha$ -CS) is a Ns-OS (resp. Ns-CS).

Proposition 2.9:[7] In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then the following statements hold and the converse of each statements are not true:

- i. Every Ng-OS (resp. Ng-CS) is a $N\alpha g$ -OS (resp. $N\alpha g$ -CS).

- ii. Every $N\alpha$ -OS (resp. $N\alpha$ -CS) is a $Ng\alpha$ -OS (resp. $Ng\alpha$ -CS).
- iii. Every $Ng\alpha$ -OS (resp. $Ng\alpha$ -CS) is a $N\alpha g$ -OS (resp. $N\alpha g$ -CS).

Proposition 2.10:[3] In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then the following statements hold and the converse of each statements are not true:

- i. Every Ng -OS (resp. Ng -CS) is a Ngs -OS (resp. Ngs -CS).
- ii. Every Ns -OS (resp. Ns -CS) is a Nsg -OS (resp. Nsg -CS).
- iii. Every Nsg -OS (resp. Nsg -CS) is a Ngs -OS (resp. Ngs -CS).
- iv. Every $Ng\alpha$ -OS (resp. $Ng\alpha$ -CS) is a Ngs -OS (resp. Ngs -CS).

3. Nano Generalized sg -Closed Sets

In this section we introduce and study the nano generalized sg -closed sets and some of its properties.

Definition 3.1: A subset \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be a nano generalized sg -closed set (briefly $Ngsg$ -CS) if $Ncl(\mathcal{A}) \subseteq \mathcal{M}$ whenever $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is a Nsg -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. The family of all $Ngsg$ -CS of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is denoted by $Ngsg-C(\mathcal{U}, X)$.

Proposition 3.2: In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, the following statements are true:

- i. Every N-CS is a $Ngsg$ -CS.
- ii. Every $Ngsg$ -CS is a Ng -CS.

Proof: (i) Let \mathcal{A} be a N-CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a Nsg -OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Then $Ncl(\mathcal{A}) = \mathcal{A} \subseteq \mathcal{M}$. Therefore \mathcal{A} is a $Ngsg$ -CS.

(ii) Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a N-OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Since every N-OS is a Nsg -OS, we have $Ncl(\mathcal{A}) \subseteq \mathcal{M}$. Therefore \mathcal{A} is a Ng -CS.

The converse of the above proposition need not be true which can be seen from the following examples.

Example 3.3: Let $\mathcal{U} = \{a, b, c, d\}$ with $\mathcal{U}/\mathcal{R} = \{\{a\}, \{d\}, \{b, c\}\}$ and $X = \{a, c\}$.

Let $\tau_{\mathcal{R}}(X) = \{\phi, \{a\}, \{b, c\}, \{a, b, c\}, \mathcal{U}\}$ be a NTS. Then the set $\{b, c\}$ is a $Ngsg$ -CS but not N-CS.

Example 3.4: Let $\mathcal{U} = \{a, b, c, d, e\}$ with $\mathcal{U}/\mathcal{R} = \{\{d\}, \{a, b\}, \{c, e\}\}$ and $X = \{a, d\}$.

Let $\tau_{\mathcal{R}}(X) = \{\phi, \{d\}, \{a, b\}, \{a, b, d\}, \mathcal{U}\}$ be a NTS. Then the set $\{a, c, d\}$ is a Ng -CS but not $Ngsg$ -CS.

Proposition 3.5: In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, the following statements are true:

- i. Every $Ngsg$ -CS is a $N\alpha g$ -CS.
- ii. Every $Ngsg$ -CS is a $Ng\alpha$ -CS.
- iii. Every $Ngsg$ -CS is a Nsg -CS.
- iv. Every $Ngsg$ -CS is a Ngs -CS.

Proof:

- i. Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a N-OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Since every N-OS is a Nsg -OS, we have $Nacl(\mathcal{A}) \subseteq Ncl(\mathcal{A}) \subseteq \mathcal{M}$ implies $Nacl(\mathcal{A}) \subseteq \mathcal{M}$. Therefore \mathcal{A} is a $N\alpha g$ -CS.
- ii. Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a $N\alpha$ -OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Since every $N\alpha$ -OS is a Ns -OS which is a Nsg -OS, we have $Nacl(\mathcal{A}) \subseteq Ncl(\mathcal{A}) \subseteq \mathcal{M}$ implies $Nacl(\mathcal{A}) \subseteq \mathcal{M}$. Therefore \mathcal{A} is a $Ng\alpha$ -CS.
- iii. Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a Ns -OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Since every Ns -OS is a Nsg -OS, we have $Nscl(\mathcal{A}) \subseteq Ncl(\mathcal{A}) \subseteq \mathcal{M}$ implies $Nscl(\mathcal{A}) \subseteq \mathcal{M}$. Therefore \mathcal{A} is a Nsg -CS.
- iv. Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be a N-OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$. Since every N-OS is a Nsg -OS, we have $Nscl(\mathcal{A}) \subseteq Ncl(\mathcal{A}) \subseteq \mathcal{M}$ implies $Nscl(\mathcal{A}) \subseteq \mathcal{M}$. Therefore \mathcal{A} is a Ngs -CS.

The converse of the above proposition need not be true as shown in the following examples.

Example 3.6: Let $\mathcal{U} = \{a, b, c, d\}$ with $\mathcal{U}/\mathcal{R} = \{\{a\}, \{c\}, \{b, d\}\}$ and $X = \{a, b\}$.

Let $\tau_{\mathcal{R}}(X) = \{\phi, \{a\}, \{b, d\}, \{a, b, d\}, \mathcal{U}\}$ be a NTS. Then the set $\{a, c\}$ is a $Ng\alpha$ -CS and hence $N\alpha g$ -CS but not $Ngsg$ -CS.

Example 3.7: Let $\mathcal{U} = \{p, q, r, s\}$ with $\mathcal{U}/\mathcal{R} = \{\{p\}, \{r\}, \{q, s\}\}$ and $X = \{p, q\}$.

Let $\tau_{\mathcal{R}}(X) = \{\phi, \{p\}, \{q, s\}, \{p, q, s\}, \mathcal{U}\}$ be a NTS. Then the set $\{p\}$ is a Nsg -CS and hence Ngs -CS but not $Ngsg$ -CS.

Remark 3.8: The $Ngsg$ -CS are independent of $N\alpha$ -CS and Ns -CS.

Definition 3.9: A subset \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be a nano generalized sg -open set (briefly $Ngsg$ -OS) iff $\mathcal{U} - \mathcal{A}$ is a $Ngsg$ -CS. The family of all $Ngsg$ -OS of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is denoted by $Ngsg-O(\mathcal{U}, X)$.

Proposition 3.10: Let $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ be a NTS. If \mathcal{A} is a N-OS, then it is a $Ngsg$ -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Let \mathcal{A} be a N-OS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $\mathcal{U} - \mathcal{A}$ is a N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. By proposition (3.2) part (i), $\mathcal{U} - \mathcal{A}$ is a $Ngsg$ -CS. Hence \mathcal{A} is a $Ngsg$ -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proposition 3.11: Let $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ be a NTS. If \mathcal{A} is a $Ngsg$ -OS, then it is a Ng -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Let \mathcal{A} be a $Ngsg$ -OS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $\mathcal{U} - \mathcal{A}$ is a $Ngsg$ -CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. By proposition (3.2) part (ii), $\mathcal{U} - \mathcal{A}$ is a Ng -CS. Hence \mathcal{A} is a Ng -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proposition 3.12: In a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, the following statements are true:

- i. Every $Ngsg$ -OS is a $N\alpha g$ -OS and $Ng\alpha$ -OS.
- ii. Every $Ngsg$ -OS is a Nsg -OS and Ng -OS.

Proof: Similar to above proposition.

Theorem 3.13: If \mathcal{A} and \mathcal{B} are $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $\mathcal{A} \cup \mathcal{B}$ is a $Ngsg$ -CS.

Proof: Let \mathcal{A} and \mathcal{B} be two $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{M} be any Nsg -OS in \mathcal{U} such that $\mathcal{A} \subseteq \mathcal{M}$ and $\mathcal{B} \subseteq \mathcal{M}$. Then we have $\mathcal{A} \cup \mathcal{B} \subseteq \mathcal{M}$. Since \mathcal{A} and \mathcal{B} are $Ngsg$ -CS in \mathcal{U} , $Ncl(\mathcal{A}) \subseteq \mathcal{M}$ and $Ncl(\mathcal{B}) \subseteq \mathcal{M}$. Now, $Ncl(\mathcal{A} \cup \mathcal{B}) = Ncl(\mathcal{A}) \cup Ncl(\mathcal{B}) \subseteq \mathcal{M}$ and so $Ncl(\mathcal{A} \cup \mathcal{B}) \subseteq \mathcal{M}$. Hence $\mathcal{A} \cup \mathcal{B}$ is a $Ngsg$ -CS in \mathcal{U} .

Theorem 3.14: If a set \mathcal{A} is $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $Ncl(\mathcal{A}) - \mathcal{A}$ contains no non-empty N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{F} be any N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ such that $\mathcal{F} \subseteq Ncl(\mathcal{A}) - \mathcal{A}$. Since \mathcal{A} is a $Ngsg$ -CS, we have $Ncl(\mathcal{A}) \subseteq \mathcal{U} - \mathcal{F}$. This implies $\mathcal{F} \subseteq \mathcal{U} - Ncl(\mathcal{A})$. Then $\mathcal{F} \subseteq Ncl(\mathcal{A}) \cap (\mathcal{U} - Ncl(\mathcal{A})) = \phi$. Thus, $\mathcal{F} = \phi$. Hence $Ncl(\mathcal{A}) - \mathcal{A}$ contains no non-empty N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Theorem 3.15: A set \mathcal{A} is $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ iff $Ncl(\mathcal{A}) - \mathcal{A}$ contains no non-empty Nsg -CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Let \mathcal{A} be a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let \mathcal{D} be any Nsg -CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ such that $\mathcal{D} \subseteq Ncl(\mathcal{A}) - \mathcal{A}$. Since \mathcal{A} is a $Ngsg$ -CS, we have $Ncl(\mathcal{A}) \subseteq \mathcal{U} - \mathcal{D}$. This implies $\mathcal{D} \subseteq \mathcal{U} - Ncl(\mathcal{A})$. Then $\mathcal{D} \subseteq Ncl(\mathcal{A}) \cap (\mathcal{U} - Ncl(\mathcal{A})) = \phi$. Thus, \mathcal{D} is empty.

Conversely, suppose that $Ncl(\mathcal{A}) - \mathcal{A}$ contains no non-empty Nsg -CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Let $\mathcal{A} \subseteq \mathcal{M}$ and \mathcal{M} is Nsg -OS. If $Ncl(\mathcal{A}) \subseteq \mathcal{M}$ then $Ncl(\mathcal{A}) \cap (\mathcal{U} - \mathcal{M})$ is non-empty. Since $Ncl(\mathcal{A})$ is N-CS and $\mathcal{U} - \mathcal{M}$ is Nsg -CS, we have $Ncl(\mathcal{A}) \cap (\mathcal{U} - \mathcal{M})$ is non-empty Nsg -CS of $Ncl(\mathcal{A}) - \mathcal{A}$ which is a contradiction. Therefore $Ncl(\mathcal{A}) \not\subseteq \mathcal{M}$. Hence \mathcal{A} is a $Ngsg$ -CS.

Theorem 3.16: If \mathcal{A} is a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and $\mathcal{A} \subseteq \mathcal{B} \subseteq Ncl(\mathcal{A})$, then \mathcal{B} is a $Ngsg$ -CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Suppose that \mathcal{A} is a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Let \mathcal{M} be a Nsg -OS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ such that $\mathcal{B} \subseteq \mathcal{M}$. Then $\mathcal{A} \subseteq \mathcal{M}$. Since \mathcal{A} is a $Ngsg$ -CS, it follows that $Ncl(\mathcal{A}) \subseteq \mathcal{M}$. Now, $\mathcal{B} \subseteq Ncl(\mathcal{A})$ implies $Ncl(\mathcal{B}) \subseteq Ncl(Ncl(\mathcal{A})) = Ncl(\mathcal{A})$. Thus, $Ncl(\mathcal{B}) \subseteq \mathcal{M}$. Hence \mathcal{B} is a $Ngsg$ -CS.

Theorem 3.17: Let $\mathcal{A} \subseteq \mathcal{V} \subseteq \mathcal{U}$ and if \mathcal{A} is a $Ngsg$ -CS in \mathcal{U} then \mathcal{A} is a $Ngsg$ -CS relative to \mathcal{V} .

Proof: $\mathcal{A} \subseteq \mathcal{V} \cap \mathcal{M}$ where \mathcal{M} is a Nsg -OS in \mathcal{U} . Then $\mathcal{A} \subseteq \mathcal{M}$ and hence $Ncl(\mathcal{A}) \subseteq \mathcal{M}$. This implies that $\mathcal{V} \cap Ncl(\mathcal{A}) \subseteq \mathcal{V} \cap \mathcal{M}$. Thus \mathcal{A} is a $Ngsg$ -CS relative to \mathcal{V} .

Proposition 3.18: If \mathcal{A} is a Nsg -OS and a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then \mathcal{A} is a N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Suppose that \mathcal{A} is a Nsg -OS and a $Ngsg$ -CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $Ncl(\mathcal{A}) \subseteq \mathcal{A}$ and since $\mathcal{A} \subseteq Ncl(\mathcal{A})$. Thus, $Ncl(\mathcal{A}) = \mathcal{A}$. Hence \mathcal{A} is a N-CS.

Theorem 3.19: For each $x \in \mathcal{U}$ either $\{x\}$ is a Nsg -CS or $\mathcal{U} - \{x\}$ is a $Ngsg$ -CS in \mathcal{U} .

Proof: If $\{x\}$ is not a Nsg -CS in \mathcal{U} then $\mathcal{U} - \{x\}$ is not a Nsg -OS and the only Nsg -OS containing $\mathcal{U} - \{x\}$ is the space \mathcal{U} itself. Therefore $Ncl(\mathcal{U} - \{x\}) \subseteq \mathcal{U}$ and so $\mathcal{U} - \{x\}$ is a $Ngsg$ -CS in \mathcal{U} .

Theorem 3.20: If \mathcal{A} and \mathcal{B} are $Ngsg$ -OS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, then $\mathcal{A} \cap \mathcal{B}$ is a $Ngsg$ -OS.

Proof: Let \mathcal{A} and \mathcal{B} be $Ngsg$ -OS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then $\mathcal{U} - \mathcal{A}$ and $\mathcal{U} - \mathcal{B}$ are $Ngsg$ -CS. By theorem (3.13), $(\mathcal{U} - \mathcal{A}) \cup (\mathcal{U} - \mathcal{B})$ is a $Ngsg$ -CS. Since $(\mathcal{U} - \mathcal{A}) \cup (\mathcal{U} - \mathcal{B}) = \mathcal{U} - (\mathcal{A} \cap \mathcal{B})$. Hence $\mathcal{A} \cap \mathcal{B}$ is a $Ngsg$ -OS.

Theorem 3.21: A set \mathcal{A} is $Ngsg$ -OS iff $\mathcal{C} \subseteq Nint(\mathcal{A})$ where \mathcal{C} is a $Ngsg$ -CS and $\mathcal{C} \subseteq \mathcal{A}$.

Proof: Suppose that $\mathcal{C} \subseteq Nint(\mathcal{A})$ where \mathcal{C} is a *Ngsg-CS* and $\mathcal{C} \subseteq \mathcal{A}$. Then $\mathcal{U} - \mathcal{A} \subseteq \mathcal{U} - \mathcal{C}$ and $\mathcal{U} - \mathcal{C}$ is a *Nsg-OS* by proposition (3.12) part (ii). Now, $Ncl(\mathcal{U} - \mathcal{A}) = \mathcal{U} - Nint(\mathcal{A}) \subseteq \mathcal{U} - \mathcal{C}$. Then $\mathcal{U} - \mathcal{A}$ is a *Ngsg-CS*. Hence \mathcal{A} is a *Ngsg-OS*.

Conversely, let \mathcal{A} be a *Ngsg-OS* and \mathcal{C} be a *Ngsg-CS* and $\mathcal{C} \subseteq \mathcal{A}$. Then $\mathcal{U} - \mathcal{A} \subseteq \mathcal{U} - \mathcal{C}$. Since $\mathcal{U} - \mathcal{A}$ is a *Ngsg-CS* and $\mathcal{U} - \mathcal{C}$ is a *Nsg-OS*, we have $Ncl(\mathcal{U} - \mathcal{A}) \subseteq \mathcal{U} - \mathcal{C}$. Then $\mathcal{C} \subseteq Nint(\mathcal{A})$.

Theorem 3.22: If $\mathcal{A} \subseteq \mathcal{B} \subseteq \mathcal{U}$ where \mathcal{A} is a *Ngsg-OS* relative to \mathcal{B} and \mathcal{B} is a *Ngsg-OS* in \mathcal{U} , then \mathcal{A} is a *Ngsg-OS* in \mathcal{U} .

Proof: Let \mathcal{F} be a *Nsg-CS* in \mathcal{U} and suppose that $\mathcal{F} \subseteq \mathcal{A}$. Then $\mathcal{F} = \mathcal{F} \cap \mathcal{B}$ is a *Nsg-CS* in \mathcal{B} . But \mathcal{A} is a *Ngsg-OS* relative to \mathcal{B} . Therefore $\mathcal{F} \subseteq Nint_{\mathcal{B}}(\mathcal{A})$. Since $Nint_{\mathcal{B}}(\mathcal{A})$ is a *N-OS* relative to \mathcal{B} . We have $\mathcal{F} \subseteq \mathcal{M} \cap \mathcal{B} \subseteq \mathcal{A}$, for some *N-OS* \mathcal{M} in \mathcal{U} . Since \mathcal{B} is a *Ngsg-OS* in \mathcal{U} , we have $\mathcal{F} \subseteq Nint(\mathcal{B}) \subseteq \mathcal{B}$. Therefore $\mathcal{F} \subseteq Nint(\mathcal{B}) \cap \mathcal{M} \subseteq \mathcal{B} \cap \mathcal{M} \subseteq \mathcal{A}$. It follows that $\mathcal{F} \subseteq Nint(\mathcal{A})$. Thus \mathcal{A} is a *Ngsg-OS* in \mathcal{U} .

Theorem 3.23: If \mathcal{A} is a *Ngsg-OS* in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and $Nint(\mathcal{A}) \subseteq \mathcal{B} \subseteq \mathcal{A}$, then \mathcal{B} is a *Ngsg-OS* in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

Proof: Suppose that \mathcal{A} is a *Ngsg-OS* in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and $Nint(\mathcal{A}) \subseteq \mathcal{B} \subseteq \mathcal{A}$. Then $\mathcal{U} - \mathcal{A}$ is a *Ngsg-CS* and $\mathcal{U} - \mathcal{A} \subseteq \mathcal{U} - \mathcal{B} \subseteq Ncl(\mathcal{U} - \mathcal{A})$. Then $\mathcal{U} - \mathcal{B}$ is a *Ngsg-CS* by theorem (3.16). Hence, \mathcal{B} is a *Ngsg-OS*.

Theorem 3.24: For a subset \mathcal{A} of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, the following statements are equivalent:

- i. \mathcal{A} is a *Ngsg-CS*.
- ii. $Ncl(\mathcal{A}) - \mathcal{A}$ contains no non-empty *Nsg-CS*.
- iii. $Ncl(\mathcal{A}) - \mathcal{A}$ is a *Ngsg-OS*.

Proof: Follows from theorem (3.15) and theorem (3.17).

Remark 3.25: The following diagram shows the relation between the different types of *N-CS*:




Diagram (3.1)

4. Nano *gsg*-Closure and Nano *gsg*-Interior

We introduce nano *gsg*-closure and nano *gsg*-interior and obtain some of its properties in this section.

Definition 4.1: The intersection of all *Ngsg-CS* in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ containing \mathcal{A} is called nano *gsg*-closure of \mathcal{A} and is denoted by $Ngsg-cl(\mathcal{A})$, $Ngsg-cl(\mathcal{A}) = \bigcap \{ \mathcal{B} : \mathcal{A} \subseteq \mathcal{B}, \mathcal{B} \text{ is a } Ngsg-CS \}$.

Definition 4.2: The union of all *Ngsg-OS* in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ contained in \mathcal{A} is called nano *gsg*-interior of \mathcal{A} and is denoted by $Ngsg-int(\mathcal{A})$, $Ngsg-int(\mathcal{A}) = \bigcup \{ \mathcal{B} : \mathcal{A} \supseteq \mathcal{B}, \mathcal{B} \text{ is a } Ngsg-OS \}$.

Proposition 4.3: Let \mathcal{A} be any set in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then the following properties hold:

- i. $Ngsg-int(\mathcal{A}) = \mathcal{A}$ iff \mathcal{A} is a *Ngsg-OS*.
- ii. $Ngsg-cl(\mathcal{A}) = \mathcal{A}$ iff \mathcal{A} is a *Ngsg-CS*.
- iii. $Ngsg-int(\mathcal{A})$ is the largest *Ngsg-OS* contained in \mathcal{A} .
- iv. $Ngsg-cl(\mathcal{A})$ is the smallest *Ngsg-CS* containing \mathcal{A} .

Proof: (i), (ii), (iii) and (iv) are obvious.

Proposition 4.4: Let \mathcal{A} be any set in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then the following properties hold:

- i. $Ngsg-int(\mathcal{U} - \mathcal{A}) = \mathcal{U} - (Ngsg-cl(\mathcal{A}))$,
- ii. $Ngsg-cl(\mathcal{U} - \mathcal{A}) = \mathcal{U} - (Ngsg-int(\mathcal{A}))$.

Proof: (i) By definition, $Ngsg-cl(\mathcal{A}) = \cap\{\mathcal{B}: \mathcal{A} \subseteq \mathcal{B}, \mathcal{B} \text{ is a } Ngsg\text{-CS}\}$

$$\begin{aligned} \mathcal{U} - (Ngsg-cl(\mathcal{A})) &= \mathcal{U} - \cap\{\mathcal{B}: \mathcal{A} \subseteq \mathcal{B}, \mathcal{B} \text{ is a } Ngsg\text{-CS}\} \\ &= \cup\{\mathcal{U} - \mathcal{B}: \mathcal{A} \subseteq \mathcal{B}, \mathcal{B} \text{ is a } Ngsg\text{-CS}\} \\ &= \cup\{\mathcal{M}: \mathcal{U} - \mathcal{A} \supseteq \mathcal{M}, \mathcal{M} \text{ is a } Ngsg\text{-OS}\} \\ &= Ngsg-int(\mathcal{U} - \mathcal{A}). \end{aligned}$$

(ii) The proof is similar to (i).

Theorem 4.5: Let \mathcal{A} and \mathcal{B} be two sets in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then the following properties hold:

- i. $Ngsg-cl(\phi) = \phi, Ngsg-cl(\mathcal{U}) = \mathcal{U}$.
- ii. $\mathcal{A} \subseteq Ngsg-cl(\mathcal{A})$.
- iii. $\mathcal{A} \subseteq \mathcal{B} \implies Ngsg-cl(\mathcal{A}) \subseteq Ngsg-cl(\mathcal{B})$.
- iv. $Ngsg-cl(\mathcal{A} \cap \mathcal{B}) \subseteq Ngsg-cl(\mathcal{A}) \cap Ngsg-cl(\mathcal{B})$.
- v. $Ngsg-cl(\mathcal{A} \cup \mathcal{B}) = Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$.
- vi. $Ngsg-cl(Ngsg-cl(\mathcal{A})) = Ngsg-cl(\mathcal{A})$.

Proof: (i) and (ii) are obvious.

(iii) By part (ii), $\mathcal{B} \subseteq Ngsg-cl(\mathcal{B})$. Since $\mathcal{A} \subseteq \mathcal{B}$, we have $\mathcal{A} \subseteq Ngsg-cl(\mathcal{B})$. But $Ngsg-cl(\mathcal{B})$ is a $Ngsg\text{-CS}$. Thus $Ngsg-cl(\mathcal{B})$ is a $Ngsg\text{-CS}$ containing \mathcal{A} . Since $Ngsg-cl(\mathcal{A})$ is the smallest $Ngsg\text{-CS}$ containing \mathcal{A} , we have $Ngsg-cl(\mathcal{A}) \subseteq Ngsg-cl(\mathcal{B})$. Hence, $\mathcal{A} \subseteq \mathcal{B} \implies Ngsg-cl(\mathcal{A}) \subseteq Ngsg-cl(\mathcal{B})$.

(iv) We know that $\mathcal{A} \cap \mathcal{B} \subseteq \mathcal{A}$ and $\mathcal{A} \cap \mathcal{B} \subseteq \mathcal{B}$. Therefore, by part (iii), $Ngsg-cl(\mathcal{A} \cap \mathcal{B}) \subseteq Ngsg-cl(\mathcal{A})$ and $Ngsg-cl(\mathcal{A} \cap \mathcal{B}) \subseteq Ngsg-cl(\mathcal{B})$. Hence $Ngsg-cl(\mathcal{A} \cap \mathcal{B}) \subseteq Ngsg-cl(\mathcal{A}) \cap Ngsg-cl(\mathcal{B})$.

(v) Since $\mathcal{A} \subseteq \mathcal{A} \cup \mathcal{B}$ and $\mathcal{B} \subseteq \mathcal{A} \cup \mathcal{B}$, it follows from part (iii) that $Ngsg-cl(\mathcal{A}) \subseteq Ngsg-cl(\mathcal{A} \cup \mathcal{B})$ and $Ngsg-cl(\mathcal{B}) \subseteq Ngsg-cl(\mathcal{A} \cup \mathcal{B})$. Hence $Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B}) \subseteq Ngsg-cl(\mathcal{A} \cup \mathcal{B})$ (1)

Since $Ngsg-cl(\mathcal{A})$ and $Ngsg-cl(\mathcal{B})$ are $Ngsg\text{-CS}$, $Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$ is also $Ngsg\text{-CS}$ by theorem (3.13). Also $\mathcal{A} \subseteq Ngsg-cl(\mathcal{A})$ and $\mathcal{B} \subseteq Ngsg-cl(\mathcal{B})$ implies that $\mathcal{A} \cup \mathcal{B} \subseteq Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$. Thus $Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$ is a $Ngsg\text{-CS}$ containing $\mathcal{A} \cup \mathcal{B}$. Since $Ngsg-cl(\mathcal{A} \cup \mathcal{B})$ is the smallest $Ngsg\text{-CS}$ containing $\mathcal{A} \cup \mathcal{B}$, we have $Ngsg-cl(\mathcal{A} \cup \mathcal{B}) \subseteq Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$ (2)

From (1) and (2), we have $Ngsg-cl(\mathcal{A} \cup \mathcal{B}) = Ngsg-cl(\mathcal{A}) \cup Ngsg-cl(\mathcal{B})$.

(vi) Since $Ngsg-cl(\mathcal{A})$ is a $Ngsg\text{-CS}$, we have by proposition (4.3) part (ii), $Ngsg-cl(Ngsg-cl(\mathcal{A})) = Ngsg-cl(\mathcal{A})$.

Theorem 4.6: Let \mathcal{A} and \mathcal{B} be two sets in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then the following properties hold:

- i. $Ngsg-int(\phi) = \phi, Ngsg-int(\mathcal{U}) = \mathcal{U}$.
- ii. $Ngsg-int(\mathcal{A}) \subseteq \mathcal{A}$.
- iii. $\mathcal{A} \subseteq \mathcal{B} \implies Ngsg-int(\mathcal{A}) \subseteq Ngsg-int(\mathcal{B})$.
- iv. $Ngsg-int(\mathcal{A} \cap \mathcal{B}) = Ngsg-int(\mathcal{A}) \cap Ngsg-int(\mathcal{B})$.
- v. $Ngsg-int(\mathcal{A} \cup \mathcal{B}) \supseteq Ngsg-int(\mathcal{A}) \cup Ngsg-int(\mathcal{B})$.
- vi. $Ngsg-int(Ngsg-int(\mathcal{A})) = Ngsg-int(\mathcal{A})$.

Proof: (i), (ii), (iii), (iv), (v) and (vi) are obvious.

Definition 4.7: A NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be a nano $T_{\frac{1}{2}}$ -space (briefly $NT_{\frac{1}{2}}$ -space) if every $Ng\text{-CS}$ in it is a $N\text{-CS}$.

Definition 4.8: A NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is said to be a nano T_{gsg} -space (briefly NT_{gsg} -space) if every $Ngsg\text{-CS}$ in it is a $N\text{-CS}$.

Proposition 4.9: Every $NT_{\frac{1}{2}}$ -space is a NT_{gsg} -space.

Proof: Let $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ be a $NT_{\frac{1}{2}}$ -space and let \mathcal{A} be a $Ngsg\text{-CS}$ in \mathcal{U} . Then \mathcal{A} is a $Ng\text{-CS}$, by proposition (3.2) part (ii). Since $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is a $NT_{\frac{1}{2}}$ -space, then \mathcal{A} is a $N\text{-CS}$ in \mathcal{U} . Hence $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is a NT_{gsg} -space.

The following example shows that the converse of the above proposition not be true.

Example 4.10: Let $\mathcal{U} = \{x, y, z\}$ with $\mathcal{U}/\mathcal{R} = \{\{x\}, \{y, z\}\}$ and $X = \{x, z\}$.

Let $\tau_{\mathcal{R}}(X) = \{\phi, \{x\}, \{y, z\}, \mathcal{U}\}$ be a NTS. Then $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is a NT_{gsg} -space but not $NT_{\frac{1}{2}}$ -space.

Theorem 4.11: For a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$, the following statements are equivalent:

- i. $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is a NT_{gsg} -space.
- ii. Every singleton of a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ is either Nsg-CS or N-OS.

Proof: (i) \Rightarrow (ii) Assume that for some $x \in \mathcal{U}$ the set $\{x\}$ is not a Nsg-CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. Then the only Nsg-OS containing $\mathcal{U} - \{x\}$ is the space \mathcal{U} itself and $\mathcal{U} - \{x\}$ is a Ngsg-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$. By assumption $\mathcal{U} - \{x\}$ is a N-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ or equivalently $\{x\}$ is a N-OS.

(ii) \Rightarrow (i) Let \mathcal{A} be a Ngsg-CS in $(\mathcal{U}, \tau_{\mathcal{R}}(X))$ and let $x \in Ncl(\mathcal{A})$. By assumption $\{x\}$ is either Nsg-CS or N-OS.

Case (1). Suppose $\{x\}$ is a Nsg-CS. If $x \notin \mathcal{A}$ then $Ncl(\mathcal{A}) - \mathcal{A}$ contains a non-empty Nsg-CS $\{x\}$ which is a contradiction to theorem (3.17). Therefore $x \in \mathcal{A}$.

Case (2). Suppose $\{x\}$ is a N-OS. Since $x \in Ncl(\mathcal{A})$, $\{x\} \cap \mathcal{A} \neq \phi$ and therefore $Ncl(\mathcal{A}) \subseteq \mathcal{A}$ or equivalently \mathcal{A} is a N-CS in a NTS $(\mathcal{U}, \tau_{\mathcal{R}}(X))$.

5. Conclusion

The class of Ngsg-CS defined using Nsg-CS forms a nano topology and lies between the class of N-CS and the class of Ng-CS. The Ngsg-CS can be used to derive a new decomposition of nano continuity and new nano separation axioms.

References

1. Lellis Thivagar, M. and Carmel Richard. **2013**. On nano forms of weakly open sets, *International Journal of Mathematics and Statistics Invention*, 1(1), pp:31-37.
2. Bhuvanewari, K. and Mythili Gnanapriya, K. **2014**. Nano generalized closed sets in nano topological spaces, *International Journal of Scientific and Research Publication*, 4(5), pp:1-3.
3. Bhuvanewari, K. and Ezhilarasi, A. **2014**. On nano semi-generalized and nano generalized-semi closed sets in nano topological spaces, *International Journal of Mathematics and Computer Applications Research*, 4(3), pp:117-124.
4. Lellis Thivagar, M. Nirmala Rebecca Paul and Saeid Jafari. **2011**. On new class of generalized closed sets, *Annals of the University of Craiova, Mathematics and Computer Science Series*, 38(3), pp:84-93.
5. Pawlak, Z. **1982**. Rough Sets, *International Journal of Information and Computer Sciences*, 11, pp:341-356.
6. Reilly, L. and Vamanamurthy. **1985**. On α -sets in topological spaces, *Tamkang Journal Math.*, 16, pp:7-11.
7. Thanga Nachiyar, R. and Bhuvanewari, K. **2014**. On nano generalized α -closed sets and nano α -generalized closed sets in nano topological spaces, *International Journal of Engineering Trends and Technology*, 13(6), pp:257-260.